

Wetland, Wildlife and Natural Features Report for Davis Subdivision in El Paso County, Colorado

June 19, 2023

Prepared for:

Bill Guman, PLA, ASLA, APA Willian Guman & Associates, Ltd. 731 North Weber Street Colorado Springs, CO 80903 Prepared by:



1455 Washburn Street Erie, Colorado 80516 (p): 970-812-3267

Project Number: 2022-22-1



TABLE OF CONTENTS

1.0 INTRODUCTION	1
1.1 Purpose	1
1.2 SITE LOCATION	1
1.3 PROJECT DESCRIPTION	2
2.0 METHODOLOGY	6
3.0 ENVIRONMENTAL SETTING	7
3.1 TOPOGRAPHY / NATURAL LANDFORM	
3.2 Soils	
3.3 VEGETATION COMMUNITIES	
3.3.1 Short- and Mixed-grass Prairie	
3.3.2 Wetland	
3.3.2 Riparian	
3.3.2 Residential Landscape	10
3.4 WETLAND HABITAT AND WATERS OF THE U.S.	
3.4.1 Methodology	12
3.4.2 Office Assessment Findings	12
3.4.3 Field Assessment Findings	
3.6 Riparian Habitat	
3.7 WILDLIFE	
4.0 FEDERAL LISTED SPECIES	
5.0 RAPTORS AND MIGRATORY BIRDS	
5.1 COGCC DATABASE	
5.2 USFWS IPAC DATA	
5.3 FIELD ASSESSMENT	
6.0 SUMMARY OF IMPACTS & RECOMMENDATIONS	
6.1 VEGETATION	
6.3 WETLAND HABITAT AND WATERS OF THE U.S.	
6.4 WILDLIFE	
6.5 FEDERAL LISTED SPECIES	
6.6 RAPTORS AND MIGRATORY BIRDS	
7.0 REGULATIONS AND RECOMMENDATIONS	
7.1 CLEAN WATER ACT	
7.2 ENDANGERED SPECIES ACT	
7.3 MIGRATORY BIRD TREATY ACT & BALD AND GOLDEN EAGLE PROTECTION	
8.0 REFERENCES	

LIST OF FIGURES

Figure 1. USGS Site Location Map	3
Figure 2. Existing Conditions Aerial Photo	4
Figure 3. Sketch Plan	5
Figure 4. Vegetation Community Map	11
Figure 5. National Wetland Inventory Map	17
Figure 6. CNHP Riparian Habitat Map	19
Figure 7. WOTUS Survey Map	20
Figure 8. Raptor Habitat Survey Map	29

LIST OF APPENDICES

APPENDIX A – USDA CUSTOM SOIL RESOURCE REPORT APPENDIX B – PHOTO LOCATION MAP AND REPRESENTATIVE PHOTOS APPENDIX C – USFWS IPAC TRUST RESOURCE REPORT APPENDIX D – USACE WETLAND DETERMINATION DATA FORMS

LIST OF ACROYNMS AND ABBREVIATIONS

AMSL	above mean sea level
BCC	Birds of Conservation Concern
BGEPA	Bald and Golden Eagle Protection Act
CDA	Colorado Department of Agriculture
CNHP	Colorado Natural Heritage Program
COGCC	Colorado Oil and Gas Conservation Commission
CPW	Colorado Parks and Wildlife
CWA	Clean Water Act
ECOS	Ecosystem Services, LLC
ESA	Endangered Species Act
Guman	Willian Guman & Associates, Ltd.
JD	jurisdictional under the Clean Water Act
LEDPA	Least Environmentally Damaging and Practicable Alternative
MBTA	Migratory Bird Treaty Act
Non-JD	non- jurisdictional under the Clean Water Act
NRCS	Natural Resource Conservation Service
NTCHS	Technical Committee for Hydric Soils
NWI	National Wetland Inventory
PCA	CNHP Potential Conservation Area
PMJM	Preble's meadow jumping mouse
Project	Davis Subdivision project
Report	Wetland, Wildlife and Natural Features Report
Site	Project site
T&E	Threatened and Endangered species
USACE	U.S. Army Corps of Engineers
USDA	U.S. Department of Agriculture
USFWS	U.S. Fish and Wildlife Service
USGS	U.S. Geological Survey

1.0 INTRODUCTION

Ecosystem Services, LLC (ECOS) was retained by Willian Guman & Associates, Ltd. (Guman) to perform a natural resource assessment for the 394.91-acre Davis Subdivision project (Project) and to prepare this Wetland, Wildlife and Natural Features Report (Report).

The contact information for the Guman and ECOS representatives for this Report is provided below:

Client

Bill Guman, PLA, ASLA, APA William Guman & Associates, Ltd. 731 North Weber Street Colorado Springs, CO 80903 Phone: (719) 633-9700 bill@guman.net

Agent

Grant E. Gurnée, P.W.S. Ecosystem Services, LLC 1455 Washburn Street Erie, Colorado 80516 Phone: (303) 746-0091 grant@ecologicalbenefits.com

1.1 Purpose

The purpose of the assessment is to compare background information with present-day conditions, ascertain the physical/ecological characteristics and conditions of the Site, identify potential environmental opportunities and constraints associated with development improvements, and determine the presence/absence and approximate extent of the following features:

- Vegetation Communities;
- Natural Landforms;
- Wetland habitat and other waters of the U.S. (i.e., lakes, ponds, streams) regulated under the Clean Water Act;
- Drainages and Riparian Areas;
- Wildlife Habitat:
 - Federal listed threatened and endangered species habitat regulated under the Endangered Species Act;
 - Migratory birds and raptors regulated under the Migratory Bird Treaty Act (MBTA) and Bald and Golden Eagle Protection Act (BEGPA).

1.2 Site Location

The Site is located approximately 2.74 miles northeast of Falcon and 4.65 miles southwest of Peyton in El Paso County, Colorado. It is situated east/southeast of Highway 24, north of Judge Orr Road, southwest of the Heritage Park subdivision, and the majority of the Site is west of Elbert Road. The Site is specifically located within the east half of the southeast quarter of Section 33 and the majority of Section 34, Township 12 South, Range 64 West in El Paso County, Colorado (El Paso County Parcels 420000354. 420000377, 4200000379, 4200000406, and 423000031). The center of the Site is located at approximately Latitude 38.961883° north, Longitude -104.543390° west at an elevation of approximately 6,780 feet above mean sea level. Refer to Figure 1, USGS Site Location Map and Figure 2, Existing Conditions Aerial Photo.

1.3 Project Description

The Applicant has developed a Sketch Plan for a combination of rural residential and commercial service uses. Please refer to Figure 3, Sketch Plan provided by the Applicant (dated May 30, 2023) and the development application for specific details and descriptions of the Project.

Figure 1

USGS Site Location Map




Figure 2 Existing Conditions Aerial Photo





Figure 3 Sketch Plan



		USE SUMMA			E MAXIMUM UNITS	2
/		LOW DENSITY RR-2.5 ZONING		32%	49	,
		LOW DENSITY RR-5 ZONING	226.30 ac	57%	43	
		COMMERCIAL CS ZONING	5.78 ac	1.5%	n.a.	
		OPEN SPACE	6.44 ac	1.7%	n.a.	
151		PARK ZONING	31.89 ac	n.a.	n.a.	
		NO-BUILD / OS	11.35 ac	2.9%	n.a.	
1		NO-BUILD	2.69 ac	1.0%	n.a.	
		TRAIL HEAD				
		15-20' BUFFER	n.a.	n.a.	n.a.	
		PROPOSED R.O.W.	20.68 ac	4.4%	n.a.	
		TOTAL: PLAIN - NO BUILD / OS' ACREAGE I ATED AS SUCH FOR EACH LOT WH			92 ACREAGE AS "PRIVATE OPEN SPAC	E" AND IS
11	BUILD /	OS.			ASEMENTS, AND PRIVATE FLOODP	
	FLOODP	ASEMENT' INCLUDES A PROPOSE LAIN - NO BUILD / OS AREAS.	D 8' WIDE COUNTY TRAIL	EASEMENT TO PROV	IDE CONNECTIVITY TO PARK AND	
20	PARCE PARCE A	L SUMMARY L ACREAGE 3.53		ED USE/ZO IMERCIAL	NE MAXIMUM U n.a.	NITS
}	B C D	13.57 7.44 31.31	RR-5 RR-2	2.5	2 DU 3 DU	
- ',	E	26.27 12.51	RR-2 RR-5 RR-2	5.0	12 DU 5 DU 5 DU	
5	G H	2.25 14.74		IMERCIAL	n.a. 5 DU	
	l J	39.59 21.85	RR-2 RR-2	2.5	16 DU 8 DU	
	K L M	28.40 69.79 57.97	RR-5 RR-5 RR-5	5.0	3 DU 14 DU 11 DU	
	N O	30.35 12.65	RR-5 RR-5	5.0	6 DU 2 DU	
	P Q	6.44 2.14	PAR		n.a. n.a.	
	RS	4.88 4.33	DET	ENTION ENTION	n.a. n.a.	
1 m	KEY	ENT PROPERTY (NAME AND ADD		KEY	NAME AND ADDR	ESS
	\bigcirc	Jane Davis Living 9060 Elbert Road Peyton, CO 80831 TSN 4200000377		(15)	Jerry F. Ager 9365 N. Curtis Roa Peyton, CO 80831 TSN 4234001001	
7	02	Junior J. Davis 9050 Elbert Road Peyton, CO 80831	- 8319	\bigcirc	Joseph A. Stenstro 9055 Elbert Road Peyton, CO 80831	om
		TSN 4200000218 Junior J. Davis 9350 Elbert Road Peyton, CO 80831	- 8319	(17)	TSN 4200000244 Thomas J. Elliott 8995 Elbert Road Peyton, CO 80831	- 8318
5	04	TSN 4200000241 Robert D. Gieck T		(18)	TSN 4200000245 Charlotte A. Howa	rd
1/1	(05)	8349 S. Allison St Littleton, CO 8012 TSN 4200000209 Mabel L. Brown		(19)	3232 Muirfield Driv Colo. Spgs., CO 80 TSN 4200000362 Brent Houser Ent.,	907
Pro 1	\bigcirc	9555 Curtis Road Peyton, CO 80831 TSN 4200000047	- 7759		11890 Garrett Roa Peyton, CO 80831 TSN 4300000539	d
7	(06)	Jennings Family 2030 Tabor Court Colorado Springs TSN 4233002007	_ `	(20) 4843	Gorilla Capital Co. 1342 High Street Eugene, OR 97401 TSN 4300000599	
) {	07	FFR7, LLC 1220 Valley Stree Colorado Springs TSN 4233002006		21)	J.D. Enghaus 14775 Judge Orr R Peyton, CO 80831	
71	08	Jennings Family 2030 Tabor Court Colorado Springs		(22) 1843	TSN 4304003001 Rodolfo Escobedo 10075 Burgess Ro Colorado Springs,	ad
		TSN 4233002004 Albert Omar Lope	Z	(23)	TSN 4304003002 Daniel Duane Com	
	\frown	7220 Rim Bluff La Colorado Springs TSN 4233002003 Roberto S. Torres	, CO 80927	(24)	8410 N. Curtis Roa Peyton, CO 80831 TSN 4304003003 Jennifer Renzelma	- 7927
	\bigcirc	9464 Winged Foo Peyton, CO 80831 TSN 4233002002	t Road	\sim	14980 Stapleton R Peyton, CO 80831 TSN 4233000034	oad
	\bigcirc	Julia B. Morgan L 4825 Old Farm Dr Colorado Springs TSN 4220000015	ive #314		George Tasoulis 227 W. Sylvestor F Littleton, CO 80129 TSN 4233000034	
/// //	\bigcirc	Maria O. Skaggs P.O. Box 219 Lake Hopatcong, TSN 4200000103	NJ 07849 - 02	(26) 19	Peter J. Hagen 704 Silver Oak Gro Colorado Springs, TSN 4200000377	
() / /	\bigcirc	Coronado Ranch 1083 Prickly Pear Colorado Springs TSN 4200000021	Place		Pete Lien & Sons, P.O. Box 440 Rapid City, SD 577 TSN 4200000405	
NIC SUN	\bigcirc	Julia B. Morgan L 4825 Old Farm Dr Colorado Springs TSN 4220000015	ive #314	28) 1089	William R. Bunn 29531 Saratoga Av Big Pine Key, FL 3 TSN 4304005019	
			0'	300'	600'	900'
11	Actu	al North				
	,		SCAL	. E: 1"	= 300' - 0	



2.0 METHODOLOGY

ECOS performed an office assessment in which available databases, resources, literature, and field guides on local flora and fauna were reviewed to gather background information on the environmental setting of the Site. We consulted several organizations, agencies, and their databases, including:

- Colorado Department of Agriculture (CDA) Noxious Weed List;
- Colorado Natural Heritage Program (CNHP);
- Colorado Oil and Gas Conservation Commission (COGCC) GIS Online;
- Colorado Parks and Wildlife (CPW);
- El Paso County Master Plan;
- El Paso County, Sub-Area Plan (provided by Client as applicable);
- Google Earth current and historic aerial imagery;
- Survey of Critical Biological Resources, El Paso County, Colorado;
- Survey of Critical Wetlands and Riparian Areas in El Paso and Pueblo Counties, Colorado;
- U.S. Army Corps of Engineers (USACE) 1987 Corps of Engineers Wetlands Delineation Manual;
- USACE 2008 Interim Regional Supplement to the Corps of Engineers Wetlands Delineation Manual: Great Plains Region;
- U.S. Department of Agriculture (USDA) PLANTS Database;
- U.S. Department of Agriculture, Natural Resources Conservation Service Soil Survey;
- U.S. Fish and Wildlife Service (USFWS) Region 6 data;
- USFWS Information, Planning, and Conservation (IPaC) database;
- USFWS National Wetland Inventory (NWI);
- U.S. Geological Survey (USGS); and
- Site-specific background data provided by Guman and their consulting Team, including topographic base mapping, site development plans, and other data pertinent to the assessment.

Following the collection and review of existing data and background information, ECOS conducted a field assessment of the Site on May 22 - 23, 2023. The purpose of the assessment was to compare background information with present-day conditions, ascertain the physical/ecological characteristics and conditions of the Site, identify potential environmental opportunities and constraints associated with development improvements, and determine the presence/absence and approximate extent of the following features:

- Vegetation Communities
- Topography / Natural Landform;
- Wetland habitat and other waters of the U.S. (i.e., lakes, ponds, streams) regulated under the Clean Water Act;
- Drainages and Riparian Areas; and

- Wildlife habitat, including:
 - Federal listed threatened and endangered species habitat regulated under the Endangered Species Act;
 - Migratory birds and raptors regulated under the Migratory Bird Treaty Act (MBTA) and Bald and Golden Eagle Protection Act (BEGPA).

During the office and on-site assessment ECOS sketched and/or mapped the above features (as applicable) with a GPS on a topographic base map provided by Guman and/or on a Google Earth aerial image of the Site. ECOS utilized GPS to document the boundaries/locations of significant natural features as deemed necessary. Representative photographs were taken to assist in describing and documenting Site conditions.

3.0 ENVIRONMENTAL SETTING

The Site is located in the Southwestern Tablelands Ecological Region (Chapman et al, 2006), which is primarily comprised of sub-humid grassland and semiarid rangeland. More specifically, the Site is located in the Foothills Grassland sub-region (26j) which contains a mix of grassland types with some small areas of isolated tallgrass prairie species that are more common much farther east. The proximity to runoff and moisture from the Front Range and the more loamy, gravelly, and deeper soils are able to support more tallgrass and midgrass species than neighboring ecoregions. Big and little bluestem and switchgrass occur, along with foothill grassland communities. The annual precipitation of 14 to 20 inches tends to be greater than in regions farther east. Soils are loamy, gravelly, moderately deep, and mesic. Rangeland and pasture are common, with small areas of cropland. Urban and suburban development has increased in recent years, expanding out from Colorado Springs and the greater Denver area.

The Site is located within the CNHP Kelso's Prairie Potential Conservation Area (PCA) according to the CNHP (CNHP, 2022), which is described as comprising B2 (Very High Biodiversity Significance) consisting of low rolling hills of tallgrass, midgrass, and shortgrass prairie with swales containing wet meadows and small ephemeral drainages that form a relatively intact landscape in north-central El Paso County. Located south and west of the Black Forest, the site encompasses the upper watershed of Black Squirrel Creek and its tributaries. Within the Kelso's Prairie site, two grassland communities have been described including one south of Highway 24 and along both sides of Judge Orr Road that includes the Davis Site. This grassland includes a fairly large occurrence of a big bluestem and little bluestem tallgrass prairie (Andropogon gerardii - Schizachyrium scoparium) which occurs in patches within about a five square mile area. The occurrence appears to be in good condition with relatively few weeds and sustainable grazing practices. Other grasses present include prairie sand reed (Calamovilfa longifolia), blue grama (Bouteloua gracilis), and scattered Indiangrass (Sorghastrum nutans). Perhaps the most striking aspect of the prairie along Judge Orr Road is the abundance of creeks and wetlands. These creeks and wetlands are supported by regional shallow groundwater resulting from groundwater recharge in the Black Forest to the north. The land gently slopes to the southeast forming the headwaters of Black Squirrel Creek. Many small

drainages flow from the area and can form wide wet meadows of up to 40 acres in size. These many drainages and wet meadows support a mosaic of wetland plants and communities including Baltic rush (*Juncus balticus* var. *montanus*), Nebraska sedge (*Carex nebrascensis*), clustered sedge (*C. praegracilis*), woolly sedge (*C. pellita*), Crawe sedge (*C. crawei*), three-square bulrush (*Scirpus pungens*), saltgrass (*Distichlis spicata*) and the European pasture grass redtop (*Agrostis gigantea*). These communities can form monotypic stands or intermingle with adjacent types.

No Critical Habitat, Wildlife Refuges, or Hatcheries are present in the vicinity of the Site according to the USFWS IPaC Trust Resources Report in Appendix B (USFWS, 2023a).

3.1 Topography / Natural Landform

The topography of the Site trends from the northwest to the southeast and is formed by three gentle ridges along the southwest, central and northeast portions of the Site, which form natural drainage depressions in the southwest, south-central and northeast portions of the Site. It ranges from a high elevation of approximately 6,818 feet above mean sea level (AMSL) along the northwest edge of the Site to a low elevation of approximately 6,720 feet AMSL in the southeastern corner of the Site.

3.2 Soils

ECOS utilized the USDA, Natural Resource Conservation Service (NRCS) Web Soil Survey (USDA, NRCS, 2023) to determine the types of soils present and if hydric soils are present within the Site, as this data assist in informing the presence/absence of potential wetland habitat regulated under the Clean Water Act. The soil data were also utilized to supplement the field observations of vegetation, as the USDA provides a correlation of native vegetation species by soil types. Please refer to the Custom Soil Resource Report for the Site in Appendix A.

The Site is comprised of the following soil types:

Map Unit Symbol & Name

- 8 Blakeland loamy sand, 1 to 9 percent slopes;
- 19 Columbine gravelly sandy loam, 0 to 3 percent slopes; and
- 29 Fluvaquentic Haploquolls, nearly level.

Pursuant to the Custom Soil Resource Report:

- The Blakeland loamy sand is not hydric; however, the 1% inclusion of Pleasant soil is hydric;
- The Columbine gravelly sandy loam is not hydric; however, the 1% inclusion of Fluvaquentic Haplaquolls and 1% inclusion of Pleasant soils are both hydric; and
- The Fluvaquentic Haplaquolls is hydric and the 1% inclusion of Haplaquolls soil is hydric as well.

Hydric soils are defined by the National Technical Committee for Hydric Soils (NTCHS, 1994) as soils that formed under conditions of saturation, flooding, or ponding long enough during the growing season to develop anaerobic conditions in the upper part. Under natural conditions, these soils are either saturated or inundated long enough during the growing season to support the growth and reproduction of hydrophytic vegetation.

If soils are wet enough for a long enough period of time to be considered hydric, they should exhibit certain properties that can be easily observed in the field. These visible properties are indicators of hydric soils. The indicators used to make onsite determinations of hydric soils are specified in *Field Indicators of Hydric Soils in the United States* (USDA, NRCS, 2010).

3.3 Vegetation Communities

3.3.1 Short- and Mixed-grass Prairie

The vegetation within the Site is primarily comprised of herbaceous short-grass prairie species with herbaceous wetland vegetation in the drainages and ephemeral swales flowing through the Site. Given the limited presence of certain mid-grass prairie species mixed throughout the shortgrass prairie, we have referred to the vegetation community as "short and mixed grass prairie". Refer to Figure 4, Vegetation Community Map. The dominant prairie grass species is blue grama (Bouteloua gracilis), with occasional little bluestem (Schizachyrium scoparium) and western wheatgrass (Pascopyrum smithii). The other most common associative prairie species are prairie aster (Machaeranthera tenacetifolia), smooth brome (Bromus inermis), fringed sage (Artemisia frigida), yucca (Yucca spp.) and prickly pear cactus (Opuntia sp.). Other species include Wood's rose (Rosa woodsii), false indigo bush (Amorpha fruticosa), sticky geranium (Geranium viscosissimum), and varrow (Achillea millefolium). The Site is moderately grazed and there are scattered weeds, including Canada thistle (Cirsium arvense), musk thistle (Carduus nutans), Scotch thistle (Onopordum acanthium), common mullein (Verbascum thapsus), horseweed (Conyza canadensis) and field bindweed (Convolvulus arvensis).

3.3.2 Wetland

Hydrophytic vegetation (wetland vegetation) is present within the northeastern, south-central, and southwest ephemeral drainages where saturated (hydric) soils are present. Dominant wetland vegetation includes Baltic rush (*Juncus balticus*), Nebraska sedge (*Carex nebrascensis*), clustered field sedged (*C. praegracilis*), saltgrass (*Distichlis spicata*), and spikerush (*Eleocharis palustris*). Dispersed sandbar willow (*Salix exigua*) is present in the northeastern ephemeral drainages. Other hydrophytic species present include water mint (*Mentha aquatica*), narrowleaf cattail (*Typha angustifolia*), and Rocky Mountain iris (*Iris missouriensis*). Refer to Figure 4, Vegetation Community Map and Figure 5, NWI Map.

3.3.2 Riparian

Riparian habitat within the Site is comprised of more robust short-grass prairie where moist, mesic soils are present adjacent to wetlands (described above) and small pockets of open water that were excavated for stock ponds (refer to Figures 4 and 6). Trees and shrubs are primarily absent, with the exception of narrowleaf and Plains cottonwood (*Populus angustifolia and deltoides*) and sandbar willow dispersed throughout the Site but mostly in the northeastern drainages. Refer to Figure 4, Vegetation Community Map and Figure 6. CNHP Riparian Habitat Map.


3.3.2 Residential Landscape

A large grove of Chinese elm (*Ulmus parvifolia*) was planted around the Davis Ranch (Gieck Ranch on USGS mapping) in the central portion of Section 34. This stand of elm serves as an excellent wind break, as well as good habitat for wildlife, including numerous bird cavities. Other common "landscape" trees such as pine, oak, and fruit trees are present in the residential areas. Refer to Figure 4, Vegetation Community Map.

Refer to Appendix B – Photo Location Map and Representative Photos of the vegetation communities found on the Site.

Figure 4

Vegetation Community Map



Source: Google Earth Aerial Image, 10/31/2022 & Ecosystem Services, LLC Site Assessment, 5/23/2023

3.4 Wetland Habitat and Waters of the U.S.

3.4.1 Methodology

ECOS utilized the USGS 7.5-minute topographic mapping, historic and current Google Earth aerial photography, the National Wetland Inventory (NWI) Wetlands Mapper (USFWS 2023). Refer to Figure 5, National Wetland Inventory Map), Colorado Wetland and Information Center – Wetlands Mapper (CNHP, 2023). Refer to Figure 6, CNHP Riparian Habitat Map and detailed Project topographic mapping (if available) to preliminarily identify potential wetland habitat and waters of the U.S. (WOTUS) on the Site. Additionally, ECOS performed a delineation with a GPS survey to identify WOTUS boundaries. Refer to Figure 7, WOTUS Survey Map.

The mapping data above was proofed during the field assessment and a formal delineation was conducted to determine the presence/absence of potential WOTUS.

The USACE wetland delineation methodology was employed to document the three field indicators (parameters) of wetland habitat (i.e., wetland hydrology, hydric soils, and a predominance of hydrophytic vegetation as explained in the *Corps of Engineers Wetlands Delineation Manual* (Environmental Laboratory, 1987) and supplemented by the *Regional Supplement to the Corps of Engineers Wetlands Delineation Manual*: Interim Regional Supplement to the Corps of Engineers of *Engineers Wetlands Delineation Manual*: Great Plains Region (USACE, 2008).

3.4.2 Office Assessment Findings

<u>USGS Mapping:</u> As referenced in *Section 3.1 Topography*, the topography of the Site trends from the northwest to the southeast and is formed by three gentle ridges along the southwest, central and northeast portions of the Site, which form three natural drainage depressions in the northeast, south-central, and southwest. USGS illustrates these drainages as follows:

- Northeastern drainage as one intermittent stream;
- South-central drainage as two intermittent branches that join into one intermittent stream; and
- Southwest drainage as one intermittent stream.

Given that the USGS Map indicates the presence of intermittent streams in all three drainages, there is a probability that they may support wetland vegetation if the sustaining hydrology is sufficient. Refer to Figure 1, USGS Site Location Map.

<u>Google Earth aerial imagery review:</u> ECOS reviewed the Site using the timelapse function in Google Earth (GE) to get a look back in time to 1985. The timeline review reveals two conspicuous drainages (the northeast and southcentral) as well as additional dry swales in the "north-central" portion of the Site, further described below:

• Northeast drainage: two branches are visible in the aerial imagery, a north and south branch:

- North branch: contains a mosaic of herbaceous, shrubby and forested wetlands. Elbert Road bisects this drainage.
- South branch: primarily a sandy wash in the upstream reach.
 A large area of potential herbaceous wetland habitat is visible downstream of the sandy wash below 2 stock pond berms.
- North-central swales: two upland swales are visible that have no apparent upstream or downstream connections to drainages and terminate in the vicinity of the two homes and out-buildings labeled on the USGS as the "Gieck Ranch". Neither swale presents a vegetation signature resembling wetland habitat.
- South-central drainage: two northern branches combine into one single drainage are visible on aerial imagery and show signatures of lush herbaceous wetland vegetation, including a vast wetland located offsite to the northwest that contributes flow into this drainage. Six on-line stock ponds are visible along this drainage. A southern branch, consisting of vast wetland complex comingles with the northern branch located mostly off-site to the west, fanning out upstream of Judge Orr Rd. The northern most tip of wetland along the south branch extends into the Site.
- Southwest drainage: Prior to June 2013 this was a sandy wash. Sometime after June 2013 and before November 2015, it appears that three cross-channel riprap/rock drainage "improvement" structures were installed across the drainage (presumably to detain water or stabilize the channel from future erosion). From 1985 to 2015 there was no vegetation signature present that may indicate herbaceous wetland habitat. Sometime after November 2015, a vegetation signature evolved that indicates herbaceous wetland habitat is present downstream of each of these riprap/rock structures. This channel flows under Stapleton Road, follows the north edge of Judge Orr Road, and ultimately drains into the South Central Drainage and associated wetlands.

No other potential wetland habitat or water bodies (natural or manmade) are evident on the aerial imagery. Refer to Figure 2, Davis Existing Conditions Aerial Photo and Figure 7, Davis Preliminary Waters and Wetlands Map.

<u>USFWS National Wetland Inventory (NWI) Wetlands Mapper:</u> The NWI Wetlands Mapper indicates the following:

- Northeast drainage:
 - North branch: This branch is identified as a potential Riverine Intermittent Streambed Seasonally Flooded (R4SBC) and Riparian Shrub-Scrub (RP1SS).
 - South branch: The downstream portion of this branch is identified as Riparian Shrub-Scrub (RP1SS). Three ponds are illustrated on this branch, two of which are identified as potential Palustrine Unconsolidated Shore Seasonally Flooded (PUSC). A patch of wetland downstream of the most downstream is identified as a potential Palustrine Emergent Persistent Temporary Flooded

(PEM1A), as it is situated in the large area of potential herbaceous wetland habitat visible in the Google Earth aerial imagery.

- North-central swales: The NWI does not identify these features as potential WOTUS.
- South-central drainage: The NWI identifies both branches as potential Palustrine Emergent Persistent Temporarily Flooded (PEM1A).
- Southwest drainage: The NWI identifies this drainage as a potential Riverine Intermittent Streambed Seasonally Flooded (R4SBC).

Refer to Figure 5, Davis National Wetland Inventory Map.

All of these drainages, except for the north-central upland swales appear to have a direct or indirect connection to Black Squirrel Creek.

<u>Colorado Wetland and Information Center – Wetlands Mapper</u>: CNHP has incorporated some of the data provided by the NWI for wetland habitat and has produced updated photo-interpretation of wetland mapping in several areas. On this Site, that data concurs with the NWI data. However, the lower reaches of the southwestern and eastern branches of the northeastern drainage are identified as Riparian Temporary Flooded Scrub-Shrub (RP1SS), a riparian habitat type that occurs adjacent to streams and is generally found on the banks of an incised channel. Refer to Figure 5, National Wetland Inventory Map and Figure 6, CNHP Riparian Habitat Map.

<u>USDA NRCS Web Soil Survey</u>: The custom soil report generated for the Site via the NRCS Web Soil Survey (USDA NRCS, 2023) identifies the presence of hydric (wetland) soil (refer to Section 3.2 and Appendix A). The USDA NRCS Soil Survey data indicate that the Fluvaquentic Haplaquolls soil type is a hydric soil and a few minor inclusions of hydric soil (1 - 2%) are components of the Blakeland and Columbine soil types. Under natural conditions, these soils are either saturated or inundated long enough during the growing season to support the growth and reproduction of hydrophytic vegetation (i.e., wetland vegetation).

3.4.3 Field Assessment Findings

The data review above and a field assessment revealed the presence of five potentially jurisdictional WOTUS features within the Site. Refer to Figure 7, WOTUS Survey Map. These five areas include:

- Northeast drainage:
 - North branch (Wetland 4)
 - South branch (Wetland 5)
- South-central drainage:
 - Northern branch (Wetlands 8 and 9)
 - Southern branch (Wetland 7)
- Southwest drainage (Wetland 6)

These natural features meet the wetland indicators and criteria that the Corps uses to assert jurisdiction pursuant to the *Corps of Engineers Wetlands Delineation Manual* (Environmental Laboratory, 1987) and associated *Interim* Regional Supplement to the Corps of Engineers Wetlands Delineation Manual: Great Plains Region (USACE, 2008). However, the final jurisdictional determination be made by the USACE.

The potentially jurisdictional WOTUS feature data is summarized below, with an explanation of the field indicators (parameters) of wetland habitat that were observed and documented by ECOS.

Northeast drainage:

North branch (Wetland 4) – The data for this branch is summarized on the W4-WET datasheet in Appendix C. The NWI labels this branch as Riverine Intermittent Streambed, Seasonally Flooded (R4SBC), and Riparian Shrub-Scrub (RP1SS); however, it is a broad, wetland swale that comprises primarily Palustrine Emergent Persistent Temporarily Flooded (PEM1A) characteristics with the inclusion of Palustrine Scrub-Shrub Broad-leaved Deciduous Temporary Flooded (PSS1A) characteristics. This feature was dominated by Baltic rush, Nebraska sedge, clustered field sedge, and sandbar willow, with inclusions of water mint, Canada thistle, and narrowleaf cattail. It is underlain by organic matter, sand, and silty clay that exhibit hydric hue and values in their matrices. Surface water was present up to approximately 1-inch depth and the soil was saturated to the surface. This area meets all 3 parameters for jurisdictional wetland habitat.

<u>South branch (Wetland 5)</u> - The data for this branch is summarized on the W5-WET datasheet in Appendix C. The NWI correctly labels this branch as PEM1A. It is a broad, wetland swale that is comprised of Palustrine Emergent vegetation including Baltic rush, Nebraska sedge, and clustered field sedge, with inclusions of Canada thistle and Rocky Mountain iris. It is underlain by silty clay loam and silty clay that exhibit hydric hue and values in their matrices. Surface water was not present and the soil was saturated at a depth of 10 inches below the surface. This area meets all 3 parameters for jurisdictional wetland habitat.

South-central drainage:

<u>North branch (Wetland 8 and 9)</u> – The data for this branch is summarized on the W8-WET and W9-WET datasheets in Appendix C. The NWI correctly identifies both branches as PEM1A. It is a broad, wetland swale.


The upstream portion is dominated by Nebraska sedge, and Baltic rush, with inclusions of water mint, narrowleaf cattail, and Canada thistle; and is underlain by organic matter, silty sand, and clay that exhibit hydric hue and values in their matrices. Surface water was present up to approximately 1-inch depth and the soil was saturated to the surface. This area meets all 3 parameters for jurisdictional wetland habitat.

The downstream portion is dominated by Baltic rush and saltgrass with clustered field sedge and is underlain by silty loam that exhibits hydric hue and values in their matrices. Surface water was not present and the soil was saturated at 12 inches below the surface. This area meets all 3 parameters for jurisdictional wetland habitat.

<u>South branch (Wetland 7)</u> – The data for this branch is summarized on the W7-WET datasheet in Appendix C. The NWI did not identify these branches, but it too is PEM1A. It is a broad, wetland swale dominated by Baltic rush and Nebraska sedge. It is underlain by sandy loam and clay that exhibit hydric hue and values in their matrices. Surface water was present up to approximately 1-inch depth and the soil was saturated at 14 inches below the surface. This area meets all 3 parameters for jurisdictional wetland habitat.

<u>Southwest drainage (Wetland 6)</u>: The data for this branch is summarized on the W6-WET datasheet in Appendix C. The NWI correctly identifies the upstream portion of this drainage as R4SBC; however, the downstream reach should be classified as PEM1A. It is a broad, wetland swale that is comprised of Baltic rush, common spikerush, and clustered field sedge, with inclusions of narrowleaf cattail. It is underlain by clayey sand that exhibits hydric hue and values in its matrices. Surface water was present up to approximately 1-inch depth and the soil was saturated to the surface. This area meets all 3 parameters for jurisdictional wetland habitat.

Figure 5 National Wetland Inventory Map



Source: Colorado Natural Heritage Program (CNHP) Wetland Mapper / U.S fish and Wildlife Service National Wetland Inventory (NWI)


3.6 Riparian Habitat

The Colorado Wetland Information Center – Wetlands Mapper (CNHP, 2023) includes the option for illustrating potential riparian habitat based on mapping produced by Colorado Parks and Wildlife (CPW). Refer to Figure 6, CNHP Riparian Habitat Map. The CPW Riparian Habitat mapping indicates the following:

- Northeast drainage (north and south):
 - The two, small, upstream ponds are identified as Unvegetated.
 - The lower reaches of the north and south branches of the northeastern drainage are identified as Open Water. The large area of potential herbaceous wetland habitat visible on Google Earth aerial imagery and identified on the NWI as emergent wetland (PEM1A) is identified as Upland Grass.
 - The trees visible in the northern branch are identified as Deciduous Cottonwood.
 - The sandy wash areas in the north and south branches are identified as Unvegetated.
- North-central swales: The CPW data does not identify these features as riparian.
- South-central drainage: The CPW identifies the upstream portion of the north branch as Herbaceous – Sedges/Rushes/Mesic Grasses (Moist Soils) Open Water. The downstream portion of the north branch is identified as Upland Vegetation which is contrary to NWI and field assessment where a vast expanse of Herbaceous – Sedges/Rushes/Mesic Grasses were found.
- Southwest drainage: The CPW data identifies this drainage as Unvegetated.


FIGURE 6

CNHP Riparian Habitat Map



Source: Colorado Natural Heritage Program (CNHP) Wetland Mapper

FIGURE 7. WOTUS Survey Map



Source: Google Earth Aerial Image, 10/31/2023 & Ecosystem Services, LLC Wetland Delineation, 5/23/2023

3.7 Wildlife

The stated purpose and intent of the "El Paso County Development Standards" wildlife section is to ensure that proposed development is reviewed with consideration of the impacts to wildlife and wildlife habitat, and to implement the provisions of the Master Plan (El Paso County, 2021). The two primary vegetation types within the Site are herbaceous prairie and wetlands. ECOS has determined that the wildlife impact potential for development of this singular Site is expected to be moderate to low, as the Site currently provides poor to moderate habitat for wildlife. Taken in a regional watershed or larger landscape context, as more and more prairie is developed over time impacts to wildlife are expected to be moderate to high as wildlife run out of space and habitat.

The Site provides habitat for prairie species such as pronghorn (*Antilocapra americana*), black-tailed prairie dog (*Cynomys ludovicianus*), thirteen-lined ground squirrel (*Ictidomys tridecemlineatus*), voles (*Microtus spp.*) and jackrabbit (*Lepus townsendii*). The Site also provides foraging and breeding habitat for predators such as coyote (*Canis latrans*), fox (*Vulpes vulpes*), badger (*Meles meles*), and occasional bobcat (*Lynx rufus*). The Site also provides good habitat for reptiles and moderate habitat for amphibians such as Woodhouse toad (*Anaxyrus woodhousii*), leopard frog (*Lithobates pipiens*), and garter snake (*Thamnophis spp.*).

The USFWS IPaC Trust Resources Report (USFWS, 2023a) (Appendix C) reports that bald eagle (*Haliaeetus leucocephalus*), golden eagle (*Aquila chrysaetos*), and ferruginous hawk (*Buteo regalis*) may utilize the area. The Site provides limited tree nesting habitat for raptors; however, ferruginous hawks may also use ground nests.

The Site contains no Critical Habitat, Wildlife Refuges, or Hatcheries according to the USFWS IPaC Trust Resources Report (USFWS, 2023a) (Appendix C).

The project proposes to develop most of the prairie, however, the drainages and grassland immediately adjacent to them within the floodplain would be preserved as Open Space. A noxious weed management plan will be implemented per State and County requirements to improve wildlife habitat, and a native plant revegetation plan for the Open Space is recommended to provide additional benefit to wildlife habitat.

4.0 FEDERAL LISTED SPECIES

A number of species that occur in El Paso County are listed as threatened and endangered (T&E) by the USFWS under the Endangered Species Act (ESA) (USFWS 2023). ECOS compiled the data regarding T&E species for the Site in Table 3 based on the Site-specific, USFWS IPaC Trust Resources Report we ran for the Project (Appendix C) and our onsite assessment. ECOS has provided our professional opinion regarding the probability that these species may occur within the Site and their probability of being impacted by the Project. The likelihood that the Project would impact any of the species listed below is insignificant to none. Most are not expected to occur in the project area and no downstream impacts are expected. The USFWS also states that there is no Critical Habitat for T&E species in the Site locations.

TABLE 3 - FEDERAL LISTED SPECIES POTENTIALLY IMPACTED BY THE PROJECT				
Species Status		Habitat Requirements and Presence	Probability of Impact by Project	
FISH				
Greenback cutthroat trout (<i>Oncorhynchus</i> <i>clarki stomias</i>)	Threatened	Cold, clear, gravely headwater streams and mountain lakes that provide an abundant food supply of insects.	None. Suitable habitat does not exist on the Site.	
Pallid sturgeon (Scaphirhynchus albus)	Endangered	Water-related activities/use in the N. Platte, S. Platte and Laramie River Basins may affect listed species in Nebraska.	None. The proposed project will not affect any of the listed river basins.	
BIRDS				
Eastern Black Rail (<i>Laterallus</i> <i>jamaicensis</i> ssp. Jamaicensis)	Threatened	Habitat includes tidally or non- tidally influenced marshes which range in salinity from salt to brackish to fresh. It requires dense overhead perennial herbaceous cover with underlying soils that are moist to saturated (occasionally dry) interspersed with or adjacent to very shallow water (typically \leq 3 cm). Eastern black rails depend on this dense cover throughout their life cycle and it is their primary strategy to avoid predation.	Insignificant. Suitable, dense, overhead, perennial, herbaceous cover and shallow water are minimal and dispersed in the discontinuous wetland habitat on the Site.	

TABLE 3 - FEDERAL LISTED SPECIES POTENTIALLY IMPACTED BY THE PROJECT				
Species	Status	Habitat Requirements and Presence	Probability of Impact by Project	
Piping plover (<i>Charadrius</i> <i>melodus</i>)	Threatened	Water-related activities/use in the N. Platte, S. Platte and Laramie River Basins may affect listed species in Nebraska.	None. The proposed project will not affect any of the listed river basins.	
MAMMALS				
Gray Wolf (<i>Canus lupis</i>)	Endangered	Inhabits a wide range of habitats including temperate forests, mountains, tundra, taiga, and grasslands. Lone, dispersing gray wolves may be present throughout the state of Colorado.	None. This species only needs to be considered if the Project activity includes a predator management program, which it does not.	

Species	Status	Habitat Requirements and Presence	Probability of Impact by Project
Preble's Meadow Jumping Mouse (Zapus hudsonius preblei)	Threatened		Very low. This species is unlikely to occur on the Site due to very limited and discontinuous riparian habitat with only a seasonal water source. No USFWS Critical Habitat is present on the Site (closest = 11.54 miles W) and no CPW Potentially Occupied Habitat is present on the Site (closest = 3.75 miles NE & 10.91 miles SW). Adjacent trapping data surrounding the Site vicinity indicates "Trapped, Not Found"; and there are no viable travel corridors from regional PMJM habitat to the Site.

TABLE 3 - FEDERAL LISTED SPECIES POTENTIALLY IMPACTED BY THE PROJECT				
Species	Status	Habitat Requirements and Presence	Probability of Impact by Project	
INSECTS				
Monarch butterfly (<i>Danaus</i> <i>plexippus</i>)	Candidate	Multigenerational migrant that breeds throughout North America and overwinters in dense congregations in Mexican montane fir forests. The larval hostplant is milkweed (<i>Asclepias</i> spp.). Habitat includes areas with nectar for feeding and/or milkweed for laying eggs, especially grasslands and wetlands. Breeding habitat threats are widespread native grassland loss and herbicide use. In Colorado, they are present in low numbers from May to September.	Insignificant. Milkweed is not present. Project impacts are undetectable relative to threats across this species' huge range. Potential impacts could be mitigated by limiting herbicide use and planting native flowering species, especially milkweed.	
PLANTS				

TABLE 3 - FEDERAL LISTED SPECIES POTENTIALLY IMPACTED BY THE PROJECT							
Species	Status	Habitat Requirements and Presence	Probability of Impact by Project				
Ute ladies'- tresses orchid (<i>Spiranthes</i> <i>diluvialis</i>)	Threatened	Primarily occurs along seasonally flooded river terraces, sub-irrigated or spring-fed abandoned stream channels or valleys, and lakeshores. May also occur along irrigation canals, berms, levees, irrigated meadows, excavated gravel pits, roadside borrow pits, reservoirs, and other human- modified wetlands.	None. Wetland areas on Site are poor quality habitat for this species and will not be impacted. The Site elevation ranges from 6,720 to 6,810 feet AMSL, which is higher than the 6,500-foot upper elevation limit documented for the species and recommended for conducting surveys by the USFWS.				

TABLE 3 - FEDERAL LISTED SPECIES POTENTIALLY IMPACTED BY THE

5.0 RAPTORS AND MIGRATORY BIRDS

Raptors and most birds are protected by the Colorado Nongame Wildlife Regulations, as well as by the federal Migratory Bird Treaty Act. Additionally, eagles are protected by the Bald and Golden Eagle Protection Act (BGEPA).

5.1 COGCC Database

ECOS utilized the Colorado Oil and Gas Conservation Commissions (COGCC) GIS Online data (<u>https://cogccmap.state.co.us/cogcc_gis_online/</u>) (COGCC, 2023) to screen the Site for potential raptor nests. No raptor nests have been mapped within one mile of the Site (COGCC, 202). The closest raptor nests to the Site are Golden Eagle and Ferruginous hawk active nests located 3.22 miles east of the eastern edge of the Site and a Golden Eagle active nest located 7.02 miles southwest of the southwest corner of the Site.

5.2 USFWS IPaC Data

The USFWS IPaC data for the Site indicates the probability of the presence of five bird species (refer to Appendix C) in the vicinity of the Site. The birds listed by IPaC are birds of particular concern either because they occur on the USFWS Birds of Conservation Concern (BCC) list or warrant special attention in the Project location. The 1988 amendment to the Fish and Wildlife Conservation Act mandates the USFWS to "identify species, subspecies, and populations of all migratory nongame birds that, without additional conservation actions, are likely to become candidates for listing under the ESA. "Birds of Conservation Concern 2021 (BCC 2021)" is the most recent effort to carry out this mandate. The birds listed by IPaC include:

- Bald Eagle (*Haliaeetus leucocephalus*) This is not a BCC in this area but warrants attention because of the BGEPA.
- Ferruginous Hawk (*Buteo regalis*) This is a BCC only in particular Bird Conservation Regions (BCRs) including Colorado. Per the USFWS Environmental Conservation Online System data (USFWS 2023b) (<u>https://ecos.fws.gov/ecp/species/6038</u>), ideal habitat for Ferruginous Hawks is grassland and shrub-steppe habitat including pastures, hayland, and cropland. Their nests can be found in trees and large shrubs and on roofs, utility structures, and artificial platforms, or near the ground on river cutbanks, or less frequently other ground locations such as rockpiles and riverbed mounds. ECOS has observed their nests open prairie habitat in this vicinity.
- Golden Eagle (Aquila chrysaetos) This is not a BCC but warrants attention because of the BGEPA
- Lesser Yellowlegs (*Tringa flavipes*) This is a BCC throughout its range in the continental USA and Alaska. Per the USFWS Per the USFWS Environmental Conservation Online System data (USFWS 2023b) (<u>https://ecos.fws.gov/ecp/species/9679</u>) the Site does not comprise suitable habitat for this species. However, they may pass through the Project vicinity in the 2nd, 3rd, and 4th week of April.


Pinyon Jay (*Gymnorhinus cyanocephalus*) - This is a BCC throughout its range in the continental USA and Alaska. Per the USFWS Environmental Conservation Online System data (USFWS 2023b) (<u>https://ecos.fws.gov/ecp/species/9420</u>) the Site does not comprise suitable habitat for this species. As their name implies, they can be found in pinyon-juniper woodland, sagebrush, scrub oak, and chaparral communities, and sometimes in pine forests. Given that Colorado is within its large Western U.S. range, this broad-brush range includes the Project vicinity, but the probability of its presence in said vicinity is limited to the 1st week in October.

5.3 Field Assessment

Two occupied Redtail hawk nests were identified in the northeastern corner of the Site. One occupied magpie nest was identified in the central portion of the Site. Two large prairie dog colonies that provide potential habitat for Burrowing owl (Athene cunicularia) are present in the south-central and southwestern portions of the Site. Numerous tree cavities were identified in live and dead trees throughout the Site, especially in the forested area around the Davis ranch. One nest was being used by a redtail hawk and one unoccupied raptor nest (likely Redtail hawk or Great Horned owl) was identified immediately south of the south Site boundary near Judge Orr Road. Numerous songbirds were detected using the Cornell Lab Merlin bird identification application, including yellow oriole (Icterus nigrogularis), red-wing blackbird (Agelaius phoeniceus), grackle (Quiscalus spp.), flicker (Colaptes spp.), meadowlark (Sturnella neglecta), Say's phoebe (Sayornis saya), western kingbird (Tyrannus verticalis), eastern kingbird (Tyrannus tyrannus), northern mocking bird (Mimus polyglottos), western woodpewee (Contopus sordidulus), house wren (Troglodytes aedon) and house sparrow (Passer domesticus). The prairie, riparian corridors, and wetland habitat may also provide nesting and foraging habitat for many other migratory birds. Please refer to Figure 8, Raptor Habitat Survey Map.

FIGURE 8.

Raptor Habitat Survey Map



Source: Google Earth Aerial Image, 10/31/2023 & Ecosystem Services, LLC MBTA Survey, 5/23/2023

6.0 SUMMARY OF IMPACTS & RECOMMENDATIONS

6.1 Vegetation

The vegetation within the Site is primarily comprised of herbaceous shortgrass prairie species. Given the presence of certain midgrass and tallgrass prairie and non-native species mixed throughout the shortgrass prairie, we have referred to the vegetation community as "short and mixed grass prairie". Wetland vegetation is comprised primarily of emergent, herbaceous, hydrophytic species in the ephemeral drainages and swales. Riparian habitat within the Site is comprised of upland grassland, herbaceous wetland, and small pockets of shallow open water (refer to Figure 6). Trees and shrubs are primarily absent, with the exception of dispersed individual narrowleaf and Plains cottonwood (*Populus angustifolia and deltoides*) and small patches of sandbar willow (*Salix exigua*) in the northeastern drainages. Refer to Figure 4, Vegetation Map.

The short and mixed grass prairie will be the primary vegetation/habitat type impacted by the proposed development. The proposed residential parcels are all planned to be low-density so that could provide ample opportunity to preserve high quality, native habitat within private lots if building envelopes/disturbance footprints are limited. Small neighborhood parks developed for tot-lots, field sports, etc. are not valuable open space for wildlife. If however they are designed to preserve some native habitat they can provide limited natural open space functions for smaller wildlife and birds. The two Commercial parcels and the internal road system are anticipated to have maximum impact on short and mixed grass prairie (e.g., 100% of the area beneath their footprint). The three Detention Ponds will result in the loss/impact primarily of short and mixed grass prairie, with minor impacts to wetland habitat resulting from stormwater outfalls into the creek systems. These impacts could be temporary and mitigated if prairie, riparian, and wetland habitat are restored after construction.

In addition to preserving the highest value existing native vegetation on public and private open space, in order to reduce overall direct impacts from the development, proposed landscaping (private and public) should consist of native prairie species from the same ecosystem that provide food and cover for wildlife. High, solid fences if proposed are a major impediment and impact wildlife movement through the landscape. Short, permeable and unbarbed wildlifefriendly fences that allow large and small species to move freely are recommended wherever fences are desired which will allow future residents to enjoy wildlife experiences in their everyday lives.

Over 80 percent of all wildlife species use riparian areas during some part of their life cycle. As such, floodplains, riparian areas including wetlands that together form linear natural corridors (i.e., greenways) should not be impacted by development and left intact. If necessary, road, trail, and utility corridors (i.e., crossings) that must cut through riparian areas should be avoided or minimized to only a few locations where the riparian corridors (and wetlands) are the narrowest or absent. Any proposed crossings should be designed perpendicular

to greenways. Greenways are ideal locations for trails that run parallel with the floodplain/riparian corridor to provide future neighborhood residents with positive natural outdoor and wildlife experiences such as bird watching (i.e., ecological benefits). The layout of the development at a sketch plan level is nebulous regarding the avoidance and minimization of impacts to greenways. During more detailed preliminary and final design, all man-made structures, including detention ponds should avoid impacting riparian areas and wetlands. Detention/water quality ponds, where required should be located adjacent to riparian areas and vegetated to the maximum extent possible utilizing native riparian and wetland vegetation in the pond bottoms; upland grasses, shrubs, and trees along side-slopes, spillways, and run-downs to expand riparian habitat for wildlife. Outfall structures from detention ponds with scour aprons are typically designed to extend into and impact wetlands and stream beds. These impacts can be mitigated by locating the outfall outside of riparian and/or wetland habitat and then creating a riparian/wetland swale that extends to the receiving stream.

Ground disturbance/removal of vegetation and exposure of soil instigates the invasion and colonization of common and noxious weeds, one of the most detrimental processes to the quality of any kind of habitat if left unchecked. As such, minimization of ground-disturbing activities that compact or remove native vegetation during construction is recommended. Thereafter, control of common, noxious weeds and non-native species in all areas (existing or landscaped) should be a priority during and after construction and as part of the long-term private residence and HOA maintenance of the Site. If native vegetation is preserved and weeds are managed, the loss of the existing habitat is minimized.

Overall impacts to vegetation communities that provide habitat for wildlife can be offset/mitigated by the thoughtful design; restrictions that minimize impacts to prairie through the employment of building envelopes; implementation of native planting and seeding requirements on private and public land; ongoing weed management; and long-term preservation of large, contiguous open space and greenways that limit crossings and fragmentation.

6.3 Wetland Habitat and Waters of the U.S.

There are five WOTUS features on the Site including the Northeast drainage (North and South branches); South-central drainage (North and South branches) and the Southwest drainage. The downstream end of the South-Central drainage collects water from the Southwest drainage and combines to form a very significant expanse of wetland along the entire north edge of Judge Orr Road. ECOS delineated the boundaries of these WOTUS features pursuant to the current USACE methodology to assist the planning and design Team in Site planning. The Sketch Plan does not reflect the locations of these delineated WOTUS features as it was prepared prior to the delineation. Therefore, during the final Site Plan design, the Project Team will incorporate avoidance and minimization of WOTUS impacts to the extent possible to meet the Least Environmentally Damaging and Practicable Alternative (LEDPA) requirements of Section 404(b)(1) of the Clean Water Act (CWA).

Based on the current Sketch Plan, Residential Parcels B, D, E, F, I, and N have WOTUS within them. WOTUS cover large portions of Parcels E and N. Commercial Parcels A and G overlap WOTUS to a lesser degree. The three Detention Pond Parcels Q, R, and S overlap WOTUS. If these ponds are designed and constructed within WOTUS, including outfall structures and scour aprons, and not held back from WOTUS boundaries, this will result in significant temporary and permanent impacts. Parcel S overlaps Wetlands 4 and 5 at the downstream ends of the north and south branches of the Northeast drainage. Parcels Q and R overlap Wetlands 8 and 10 at the upstream and downstream ends of the north branch of the South-Central drainage. The internal Residential Collector road system as currently laid out will have a "crossing" impact on Wetlands 7 and 8 along the north and south branch of the South-Central drainage unless Wetland 8 is free-spanned by a bridge and this road rerouted around Wetland 7. It is highly likely that "drainage improvements" like drop or grade control structures will be required by the County to decrease velocity and shear stress in all the major drainages on the Site which will result in additional impacts to WOTUS. Refer to Figure 3, Sketch Plan and Figure 7, WOTUS Survey Map.

If the impacts remain as proposed in the current Sketch Plan, the Project will require a CWA Section 404 permit. The specific type of permit cannot be identified until the final Site Plan is complete and final impacts are assessed. ECOS will work with the planning and design Team to assist in incorporating avoidance and minimization of WOTUS impacts during subsequent planning and design phases of the Project.

6.4 Wildlife

The impact to wildlife would be similar to that for vegetation. Elimination of grassland areas (native or non-native alike) and reduction of open space would have an overall negative and landscape-scale impact on wildlife species as is the case with all development spreading out over plains. The highest quality habitats (i.e., floodplains, riparian areas, and wetlands within each of the drainage systems) on the Site should be preserved as contiguous open space to help meet the life requisites of wildlife. Native grassland on private lots will be the most impacted by development and therefore efforts should be made to limit development to specific building envelopes. Weedy grassland should be managed to restore their health to improve their functional capacity to provide food, cover, and breeding habitat for all obligate prairie species that typically utilize grasslands to meet their life needs. Native landscaping around all residential and commercial structures can benefit wildlife, especially small wildlife including insects, rodents, and birds. Upland, riparian, and wetland habitat may be enhanced or created within and adjacent to a proposed detention/stormwater quality detention basins to expand upon existing riparian greenways. Management priorities should include weed control and enhancement of existing native vegetation throughout the entire development, including preserved floodplains. Altogether, a low-impact development approach that preserves grassland on private and public land combined with vigilant management actions

to maintain it will help mitigate the negative impacts to wildlife communities at a landscape scale.

6.5 Federal Listed Species

The Site is not located within any officially designated occupied or critical habitat for federally designated T&E species. Therefore, there will be no impacts to federally designated T&E species and no need to initiate consultation with the USFWS under the ESA.

6.6 Raptors and Migratory Birds

The Project is expected to have a slightly negative impact on raptors and migratory birds since open space, grassland, and hunting grounds will be lost to development. Potential habitat for burrowing owls (i.e., prairie dog colonies) and many other species that rely upon prairie dogs as a keystone species for food and their burrows for shelter will diminish their viability as species over time as more and more of the prairie is developed. Preservation of high-value wetlands and riparian areas and upland areas within the floodplain along drainages and integration of native prairie and native plantings within the fabric of the development would partially mitigate the loss of prairie, but not for burrowing owl.

7.0 REGULATIONS AND RECOMMENDATIONS

7.1 Clean Water Act

Section 404 of the CWA prohibits the discharge of dredged or fill material into WOTUS (including wetland habitat) without a valid 404 permit. ECOS identified potentially jurisdictional WOTUS (drainages with a defined bed and bank and/or persistent, abutting, connected, and continuous wetlands) that will likely require a 404 permit for any proposed impacts. However, given the current, actively changing regulatory environment at the Federal level (i.e., revision of the definition of WOTUS via the Sackett vs. USEPA Supreme Court decision) it is not feasible to determine with certainty which drainages will be deemed jurisdictional by the USACE without going through a formal jurisdictional determination process. In addition, the state of Colorado is developing a regulatory framework to protect and regulate waters and wetland of the State as a means to accommodate the WOTUS features that may be excluded from federal jurisdiction.

Floodplains, riparian areas, wetlands, and streams provide numerous cultural, ecological, and economic functions and values for society, including food and habitat for fish and wildlife, water quality improvement; flood storage; erosion control; economically beneficial natural products for human use; open space for recreation and education; and views and aesthetic qualities that improve real estate sales and values. Regardless of jurisdictional status, the waterways and wetlands present on site should be preserved to achieve these functions and values.

7.2 Endangered Species Act

The Site is not located within any officially designated occupied or critical habitat for federally designated threatened or endangered species, including the Preble's meadow jumping mouse. Therefore, there will be no impacts to federally designated threatened or endangered species and no need to initiate consultation with the USFWS under the ESA.

Please note the following standard response from the USFWS in regard to ESA concurrence or clearance: "If you (the project proponent) have determined that your project will have no effect to listed species or their habitat, or if suitable habitat for a listed species does not occur within your project area, you may not receive any further response or notification from us, as neither section 7 of the Endangered Species Act of 1973, as amended (ESA; 16 U.S.C 1531 *et seq.*), nor implementing regulations under section 7 of the ESA, require us to review or concur with projects where "no effect" determinations have been made". This means that the USFWS may or may not comment or provide effects determinations as documentation of ESA compliance regardless of the Project being constructed, funded or permitted by a federal agency or if requested by the County or FEMA.

7.3 Migratory Bird Treaty Act & Bald and Golden Eagle Protection Act

No raptor nests have been mapped within one mile outside of the Site (COGCC 2023. However, migratory bird and raptor nests, and expansive prairie dog colonies that are potential habitat for burrowing owls were observed within the Site (refer to Section 5.3). Given the seasonal and transitory nature of migratory birds and raptors, including burrowing owls, ECOS recommends a nesting bird survey immediately prior to any construction activity to identify any new or active nests or burrows within the Site or within the CPW recommended buffers of the Site. Construction activities should be restricted during the breeding season near any identified active migratory bird nest or burrow.

8.0 REFERENCES

COGCC (Colorado Oil and Gas Conservation Commission). 2023. COGCC GIS Online. Available at: <u>https://cogccmap.state.co.us/cogcc_gis_online/</u>

CNHP. 2001. Survey of Critical Wetlands and Riparian Areas in El Paso and Pueblo Counties, Colorado. Colorado Natural Heritage Program, College of Natural Resources, 254 General Services Building, Colorado State University, Fort Collins, CO 80523. June 27, 2001.

CNHP. 2023. Colorado Wetland Inventory, Wetlands Mapper. Available at: <u>https://csurams.maps.arcgis.com/apps/webappviewer/index.html</u>.

Environmental Laboratory. 1987. Corps of Engineers Wetlands Delineation Manual. Technical Report Y-87-1, U.S. Army Engineer Waterways Experiment Station, Vicksburg, Mississippi.

Kershaw, Linda, A. MacKinnon, and J. Pojar. 1998. Plants of the Rocky Mountains. Lone Pine Publishing, Edmonton, Canada.

NTCHS (National Technical Committee for Hydric Soils). 1994. *Changes in Hydric Soils of the United States* (including the NTCHS definition of Hydric Soil). Federal Register Volume 59, Number 133. Wednesday, July 13, 1994.

USACE (U.S. Army Corps of Engineers). 2008. Regional Supplement to the Corps of Engineers Wetlands Delineation Manual: Interim Regional Supplement to the Corps of Engineers Wetlands Delineation Manual: Great Plains Region. May 2008.

USDA (U.S. Department of Agriculture). 2023. USDA PLANTS Database. Available at: <u>http://plants.usda.gov/java/</u>.

USDA, Natural Resources Conservation Service (NRCS). 2010. Field Indicators of Hydric Soils in the United States, A Guide for Identifying and Delineating Hydric Soils, Version 7.0. L.M. Vasilas, G.W. Hurt and C.V. Noble (eds.). USDA, NRCS, in cooperation with the National Technical Committee for Hydric Soils.

USDA, NRCS. 2015. National Hydric Soils List for Colorado. Available at: <u>https://www.codot.gov/programs/environmental/wetlands/tools.html</u> and select the Colorado Hydric Soils List in the drop-down menu.

USDA, NRCS. 2023. Web Soil Survey. Available at: <u>http://websoilsurvey.nrcs.usda.gov/app/WebSoilSurvey.aspx</u>.

USFWS (U.S. Fish and Wildlife Service). 2023a. Information, Planning, and Conservation System. Available at: <u>http://ecos.fws.gov/ipac/</u>.

USFWS. 2023b. Environmental Conservation Online System. Available at: http://ecos.fws.gov/tess_public/countySearch!speciesByCountyReport.action?fip_s=08049.

Weber, William A. and R.C. Wittmann. 2012. Colorado Flora: Eastern Slope, Fourth Edition. University Press of Colorado, Boulder, Colorado.

Whitson, Tom D. L.C. Burrill, S.A. Dewey, D.W. Cudney, B.E. Nelson, R.D. Lee, and R. Parker. 2004. Weeds of the West, 9th Edition. Western Society of Weed Science, Western United States Land Grant Universities Cooperative Extension Services, and the University of Wyoming, Jackson Hole, Wyoming.

Wingate, Janet. L. 1994. Illustrated Key to the Grasses of Colorado. Wingate Consulting, Denver, Colorado.

Appendix A

USDA Soil Data



United States Department of Agriculture


Natural Resources Conservation

Service

A product of the National Cooperative Soil Survey, a joint effort of the United States Department of Agriculture and other Federal agencies, State agencies including the Agricultural Experiment Stations, and local participants

Custom Soil Resource Report for El Paso County Area, Colorado

Davis Site



Preface

Soil surveys contain information that affects land use planning in survey areas. They highlight soil limitations that affect various land uses and provide information about the properties of the soils in the survey areas. Soil surveys are designed for many different users, including farmers, ranchers, foresters, agronomists, urban planners, community officials, engineers, developers, builders, and home buyers. Also, conservationists, teachers, students, and specialists in recreation, waste disposal, and pollution control can use the surveys to help them understand, protect, or enhance the environment.

Various land use regulations of Federal, State, and local governments may impose special restrictions on land use or land treatment. Soil surveys identify soil properties that are used in making various land use or land treatment decisions. The information is intended to help the land users identify and reduce the effects of soil limitations on various land uses. The landowner or user is responsible for identifying and complying with existing laws and regulations.

Although soil survey information can be used for general farm, local, and wider area planning, onsite investigation is needed to supplement this information in some cases. Examples include soil quality assessments (http://www.nrcs.usda.gov/wps/portal/nrcs/main/soils/health/) and certain conservation and engineering applications. For more detailed information, contact your local USDA Service Center (https://offices.sc.egov.usda.gov/locator/app?agency=nrcs) or your NRCS State Soil Scientist (http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/contactus/? cid=nrcs142p2_053951).

Great differences in soil properties can occur within short distances. Some soils are seasonally wet or subject to flooding. Some are too unstable to be used as a foundation for buildings or roads. Clayey or wet soils are poorly suited to use as septic tank absorption fields. A high water table makes a soil poorly suited to basements or underground installations.

The National Cooperative Soil Survey is a joint effort of the United States Department of Agriculture and other Federal agencies, State agencies including the Agricultural Experiment Stations, and local agencies. The Natural Resources Conservation Service (NRCS) has leadership for the Federal part of the National Cooperative Soil Survey.

Information about soils is updated periodically. Updated information is available through the NRCS Web Soil Survey, the site for official soil survey information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or a part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require


alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410 or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.

Contents

Preface	2
Soil Map	
Soil Map	
Legend	
Map Unit Legend	
Map Unit Descriptions	
El Paso County Area, Colorado	
8—Blakeland loamy sand, 1 to 9 percent slopes	10
19—Columbine gravelly sandy loam, 0 to 3 percent slopes	11
29—Fluvaquentic Haplaquolls, nearly level	12
References	14

Soil Map

The soil map section includes the soil map for the defined area of interest, a list of soil map units on the map and extent of each map unit, and cartographic symbols displayed on the map. Also presented are various metadata about data used to produce the map, and a description of each soil map unit.



MAP LEGEND)	MAP INFORMATION		
Area of In	terest (AOI) Area of Interest (AOI)	00	Spoil Area	The soil surveys that comprise your AOI were mapped at 1:24,000.		
	Area of Interest (AOI)	۵	Stony Spot			
Soils	Soil Map Unit Polygons	0	Very Stony Spot	Warning: Soil Map may not be valid at this scale.		
~	Soil Map Unit Lines	\$	Wet Spot			
	Soil Map Unit Points	\triangle	Other	Enlargement of maps beyond the scale of mapping can cause misunderstanding of the detail of mapping and accuracy of soil		
		Special Line Features Water Features		line placement. The maps do not show the small areas of		
Special Point Features Blowout				contrasting soils that could have been shown at a more detailed scale.		
×	Borrow Pit	\sim	Streams and Canals			
×	Clay Spot	Transport		Please rely on the bar scale on each map sheet for map		
~	Closed Depression	+++	Rails	measurements.		
~	Gravel Pit	~	Interstate Highways	Source of Map: Natural Resources Conservation Service		
X		~	US Routes	Web Soil Survey URL: Coordinate System: Web Mercator (EPSG:3857)		
	Gravelly Spot	~	Major Roads			
٥	Landfill	\sim	Local Roads	Maps from the Web Soil Survey are based on the Web Mercator		
٨.	Lava Flow	Backgrou		projection, which preserves direction and shape but distorts distance and area. A projection that preserves area, such as the		
عليه	Marsh or swamp	Mar.	Aerial Photography	Albers equal-area conic projection, should be used if more		
∞	Mine or Quarry			accurate calculations of distance or area are required.		
0	Miscellaneous Water			This product is generated from the USDA-NRCS certified data a of the version date(s) listed below.		
0	Perennial Water					
\vee	Rock Outcrop			Soil Survey Area: El Paso County Area, Colorado		
+	Saline Spot			Survey Area Data: Version 20, Sep 2, 2022		
°.°	Sandy Spot			Soil map units are labeled (as space allows) for map scales		
-	Severely Eroded Spot			1:50,000 or larger.		
\diamond	Sinkhole			Date(s) aerial images were photographed: Sep 11, 2018—Oct		
3	Slide or Slip			20, 2018		
ø	Sodic Spot			The orthophoto or other base map on which the soil lines were compiled and digitized probably differs from the background imagery displayed on these maps. As a result, some minor shifting of map unit boundaries may be evident.		

Map Unit Legend

Map Unit Symbol	Map Unit Name	Acres in AOI	Percent of AOI	
8	Blakeland loamy sand, 1 to 9 percent slopes	109.7	28.1%	
19	Columbine gravelly sandy loam, 0 to 3 percent slopes	243.5	62.3%	
29	Fluvaquentic Haplaquolls, nearly level	37.4	9.6%	
Totals for Area of Interest		390.5	100.0%	

Map Unit Descriptions

The map units delineated on the detailed soil maps in a soil survey represent the soils or miscellaneous areas in the survey area. The map unit descriptions, along with the maps, can be used to determine the composition and properties of a unit.

A map unit delineation on a soil map represents an area dominated by one or more major kinds of soil or miscellaneous areas. A map unit is identified and named according to the taxonomic classification of the dominant soils. Within a taxonomic class there are precisely defined limits for the properties of the soils. On the landscape, however, the soils are natural phenomena, and they have the characteristic variability of all natural phenomena. Thus, the range of some observed properties may extend beyond the limits defined for a taxonomic class. Areas of soils of a single taxonomic class rarely, if ever, can be mapped without including areas of other taxonomic classes. Consequently, every map unit is made up of the soils or miscellaneous areas for which it is named and some minor components that belong to taxonomic classes other than those of the major soils.

Most minor soils have properties similar to those of the dominant soil or soils in the map unit, and thus they do not affect use and management. These are called noncontrasting, or similar, components. They may or may not be mentioned in a particular map unit description. Other minor components, however, have properties and behavioral characteristics divergent enough to affect use or to require different management. These are called contrasting, or dissimilar, components. They generally are in small areas and could not be mapped separately because of the scale used. Some small areas of strongly contrasting soils or miscellaneous areas are identified by a special symbol on the maps. If included in the database for a given area, the contrasting minor components are identified in the map unit descriptions along with some characteristics of each. A few areas of minor components may not have been observed, and consequently they are not mentioned in the descriptions, especially where the pattern was so complex that it was impractical to make enough observations to identify all the soils and miscellaneous areas on the landscape.

The presence of minor components in a map unit in no way diminishes the usefulness or accuracy of the data. The objective of mapping is not to delineate pure taxonomic classes but rather to separate the landscape into landforms or

landform segments that have similar use and management requirements. The delineation of such segments on the map provides sufficient information for the development of resource plans. If intensive use of small areas is planned, however, onsite investigation is needed to define and locate the soils and miscellaneous areas.

An identifying symbol precedes the map unit name in the map unit descriptions. Each description includes general facts about the unit and gives important soil properties and qualities.

Soils that have profiles that are almost alike make up a *soil series*. Except for differences in texture of the surface layer, all the soils of a series have major horizons that are similar in composition, thickness, and arrangement.

Soils of one series can differ in texture of the surface layer, slope, stoniness, salinity, degree of erosion, and other characteristics that affect their use. On the basis of such differences, a soil series is divided into *soil phases*. Most of the areas shown on the detailed soil maps are phases of soil series. The name of a soil phase commonly indicates a feature that affects use or management. For example, Alpha silt loam, 0 to 2 percent slopes, is a phase of the Alpha series.

Some map units are made up of two or more major soils or miscellaneous areas. These map units are complexes, associations, or undifferentiated groups.

A *complex* consists of two or more soils or miscellaneous areas in such an intricate pattern or in such small areas that they cannot be shown separately on the maps. The pattern and proportion of the soils or miscellaneous areas are somewhat similar in all areas. Alpha-Beta complex, 0 to 6 percent slopes, is an example.

An *association* is made up of two or more geographically associated soils or miscellaneous areas that are shown as one unit on the maps. Because of present or anticipated uses of the map units in the survey area, it was not considered practical or necessary to map the soils or miscellaneous areas separately. The pattern and relative proportion of the soils or miscellaneous areas are somewhat similar. Alpha-Beta association, 0 to 2 percent slopes, is an example.

An *undifferentiated group* is made up of two or more soils or miscellaneous areas that could be mapped individually but are mapped as one unit because similar interpretations can be made for use and management. The pattern and proportion of the soils or miscellaneous areas in a mapped area are not uniform. An area can be made up of only one of the major soils or miscellaneous areas, or it can be made up of all of them. Alpha and Beta soils, 0 to 2 percent slopes, is an example.

Some surveys include *miscellaneous areas*. Such areas have little or no soil material and support little or no vegetation. Rock outcrop is an example.

El Paso County Area, Colorado

8-Blakeland loamy sand, 1 to 9 percent slopes

Map Unit Setting

National map unit symbol: 369v Elevation: 4,600 to 5,800 feet Mean annual precipitation: 14 to 16 inches Mean annual air temperature: 46 to 48 degrees F Frost-free period: 125 to 145 days Farmland classification: Not prime farmland

Map Unit Composition

Blakeland and similar soils: 98 percent Minor components: 2 percent Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Blakeland

Setting

Landform: Flats, hills Landform position (three-dimensional): Side slope, talf Down-slope shape: Linear Across-slope shape: Linear Parent material: Alluvium derived from sedimentary rock and/or eolian deposits derived from sedimentary rock

Typical profile

A - 0 to 11 inches: loamy sand AC - 11 to 27 inches: loamy sand C - 27 to 60 inches: sand

Properties and qualities

Slope: 1 to 9 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Somewhat excessively drained
Runoff class: Low
Capacity of the most limiting layer to transmit water (Ksat): High to very high (5.95 to 19.98 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Calcium carbonate, maximum content: 5 percent
Available water supply, 0 to 60 inches: Low (about 4.5 inches)

Interpretive groups

Land capability classification (irrigated): 3e Land capability classification (nonirrigated): 6e Hydrologic Soil Group: A Ecological site: R049XB210CO - Sandy Foothill Hydric soil rating: No

Minor Components

Other soils

Percent of map unit: 1 percent

Hydric soil rating: No

Pleasant

Percent of map unit: 1 percent Landform: Depressions Hydric soil rating: Yes

19—Columbine gravelly sandy loam, 0 to 3 percent slopes

Map Unit Setting

National map unit symbol: 367p Elevation: 6,500 to 7,300 feet Mean annual precipitation: 14 to 16 inches Mean annual air temperature: 46 to 50 degrees F Frost-free period: 125 to 145 days Farmland classification: Not prime farmland

Map Unit Composition

Columbine and similar soils: 97 percent Minor components: 3 percent Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Columbine

Setting

Landform: Fans, fan terraces, flood plains Down-slope shape: Linear Across-slope shape: Linear Parent material: Alluvium

Typical profile

A - 0 to 14 inches: gravelly sandy loam *C - 14 to 60 inches:* very gravelly loamy sand

Properties and qualities

Slope: 0 to 3 percent
Depth to restrictive feature: More than 80 inches
Drainage class: Well drained
Runoff class: Very low
Capacity of the most limiting layer to transmit water (Ksat): High to very high (5.95 to 19.98 in/hr)
Depth to water table: More than 80 inches
Frequency of flooding: None
Frequency of ponding: None
Available water supply, 0 to 60 inches: Very low (about 2.5 inches)

Interpretive groups

Land capability classification (irrigated): 4e Land capability classification (nonirrigated): 6e Hydrologic Soil Group: A Ecological site: R049XY214CO - Gravelly Foothill Hydric soil rating: No

Minor Components

Fluvaquentic haplaquolls

Percent of map unit: 1 percent Landform: Swales Hydric soil rating: Yes

Other soils

Percent of map unit: 1 percent *Hydric soil rating:* No

Pleasant

Percent of map unit: 1 percent Landform: Depressions Hydric soil rating: Yes

29—Fluvaquentic Haplaquolls, nearly level

Map Unit Setting

National map unit symbol: 3681 Elevation: 5,000 to 7,800 feet Mean annual precipitation: 13 to 15 inches Mean annual air temperature: 46 to 52 degrees F Frost-free period: 110 to 165 days Farmland classification: Not prime farmland

Map Unit Composition

Fluvaquentic haplaquolls and similar soils: 98 percent *Minor components:* 2 percent *Estimates are based on observations, descriptions, and transects of the mapunit.*

Description of Fluvaquentic Haplaquolls

Setting

Landform: Marshes, flood plains, swales Down-slope shape: Linear Across-slope shape: Linear Parent material: Alluvium

Typical profile

A - 0 to 12 inches: variable

C - 12 to 60 inches: stratified very gravelly sand to loam

Properties and qualities

Slope: 0 to 2 percent Depth to restrictive feature: More than 80 inches Drainage class: Poorly drained Runoff class: Very high Capacity of the most limiting layer to transmit water (Ksat): Moderately high to high (0.20 to 6.00 in/hr) Depth to water table: About 0 to 24 inches Frequency of flooding: Frequent Frequency of ponding: None Maximum salinity: Nonsaline to slightly saline (0.0 to 4.0 mmhos/cm) Available water supply, 0 to 60 inches: Moderate (about 6.2 inches)

Interpretive groups

Land capability classification (irrigated): None specified Land capability classification (nonirrigated): 5w Hydrologic Soil Group: D Ecological site: R067BY029CO - Sandy Meadow Hydric soil rating: Yes

Minor Components

Haplaquolls

Percent of map unit: 1 percent Landform: Domes Hydric soil rating: Yes

Other soils

Percent of map unit: 1 percent Hydric soil rating: No

References

American Association of State Highway and Transportation Officials (AASHTO). 2004. Standard specifications for transportation materials and methods of sampling and testing. 24th edition.

American Society for Testing and Materials (ASTM). 2005. Standard classification of soils for engineering purposes. ASTM Standard D2487-00.

Cowardin, L.M., V. Carter, F.C. Golet, and E.T. LaRoe. 1979. Classification of wetlands and deep-water habitats of the United States. U.S. Fish and Wildlife Service FWS/OBS-79/31.

Federal Register. July 13, 1994. Changes in hydric soils of the United States.

Federal Register. September 18, 2002. Hydric soils of the United States.

Hurt, G.W., and L.M. Vasilas, editors. Version 6.0, 2006. Field indicators of hydric soils in the United States.

National Research Council. 1995. Wetlands: Characteristics and boundaries.

Soil Survey Division Staff. 1993. Soil survey manual. Soil Conservation Service. U.S. Department of Agriculture Handbook 18. http://www.nrcs.usda.gov/wps/portal/ nrcs/detail/national/soils/?cid=nrcs142p2_054262

Soil Survey Staff. 1999. Soil taxonomy: A basic system of soil classification for making and interpreting soil surveys. 2nd edition. Natural Resources Conservation Service, U.S. Department of Agriculture Handbook 436. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2_053577

Soil Survey Staff. 2010. Keys to soil taxonomy. 11th edition. U.S. Department of Agriculture, Natural Resources Conservation Service. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2_053580

Tiner, R.W., Jr. 1985. Wetlands of Delaware. U.S. Fish and Wildlife Service and Delaware Department of Natural Resources and Environmental Control, Wetlands Section.

United States Army Corps of Engineers, Environmental Laboratory. 1987. Corps of Engineers wetlands delineation manual. Waterways Experiment Station Technical Report Y-87-1.

United States Department of Agriculture, Natural Resources Conservation Service. National forestry manual. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/ home/?cid=nrcs142p2 053374

United States Department of Agriculture, Natural Resources Conservation Service. National range and pasture handbook. http://www.nrcs.usda.gov/wps/portal/nrcs/ detail/national/landuse/rangepasture/?cid=stelprdb1043084

United States Department of Agriculture, Natural Resources Conservation Service. National soil survey handbook, title 430-VI. http://www.nrcs.usda.gov/wps/portal/ nrcs/detail/soils/scientists/?cid=nrcs142p2_054242

United States Department of Agriculture, Natural Resources Conservation Service. 2006. Land resource regions and major land resource areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/? cid=nrcs142p2_053624

United States Department of Agriculture, Soil Conservation Service. 1961. Land capability classification. U.S. Department of Agriculture Handbook 210. http://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/nrcs142p2_052290.pdf

Appendix B

Photo Location Map and Representative Photos

PHOTO LOCATION MAP (Representative photos taken on 5/23/2023)






Photo Point REP 1 – Photo looking south of Short- and Mixed-grass Prairie and prairie dog colony on the parcel immediately west of Stapleton Road.



Photo Point REP 2 – Photo looking north of Short- and Mixed-grass Prairie and prairie dog colony on the portion of the site immediately east of Stapleton Road.



Photo Point REP 2 – Photo looking east of Short- and Mixed-grass Prairie and prairie dog colony on the portion of the site immediately east of Stapleton Road.



Photo Point REP 3 – Photo looking southeast of Short- and Mixed-grass Prairie immediately south of the Davis ranch.



Photo Point REP 3 – Photo looking south of Short- and Mixed-grass Prairie immediately south of the Davis ranch.



Photo Point REP 3 – Photo looking southwest of Short- and Mixed-grass Prairie immediately south of the Davis ranch.



Photo Point REP 3 – Photo looking northwest of Residential Landscape surrounding the Davis ranch.



Photo Point REP 3 – Photo looking north of Residential Landscape surrounding the Davis ranch.



Photo Point REP 3 – Photo looking north of Residential Landscape surrounding the Davis ranch.



Photo Point REP 4 – Photo looking west of Residential Landscape (Chinese elm) grove surrounding the Davis ranch.



Photo Point REP 5 – Photo looking northwest of Short- and Mixed-grass Prairie immediately north of the Davis ranch.



Photo Point REP 5 – Photo looking north of Short- and Mixed-grass Prairie immediately north of the Davis ranch.



Photo Point REP 5 – Photo looking northeast of Short- and Mixed-grass Prairie immediately north of the Davis ranch.



Photo Point REP 5 – Photo looking east of Short- and Mixed-grass Prairie, junkyard and Chinese elm grove immediately north of the Davis ranch.



Photo Point W4 WET – Photo looking northwest of riparian and wetland habitat in the north branch of the Northeast drainage (Wetland 4).



Photo Point W4 WET – Photo looking southeast of riparian and wetland habitat in the north branch of the Northeast drainage (Wetland 4) near Elbert Road.



Photo Point W5 WET – Photo looking northwest of riparian and wetland habitat in the south branch of the Northeast drainage (Wetland 5).



Photo Point W5 WET – Photo looking southeast of riparian and wetland habitat in the south branch of the Northeast drainage (Wetland 5) near Elbert Road.



Photo Point W6 WET – Photo looking southeast of wetland habitat between drop structures in Southwest drainage (Wetland 6).



Photo Point W7 WET – Photo looking north of upper end of groundwater fed wetland (Wetland 7) within the south branch of the South-central drainage.



Photo Point W7 WET – Photo looking south of ground water fed wetland complex (Wetland 7) within the south branch of the South-central drainage.



Photo Point W8 WET US – Photo looking northwest (upstream/off-site) of wetland habitat in the north branch of the South-central drainage (Wetland 8).



Photo Point W8 WET DS – Photo looking southeast (downstream/on-site) of wetland habitat in the north branch of the South-central drainage (Wetland 8).



Photo Point W8 WET DS – Photo looking northwest (downstream/on-site) of riparian and wetland habitat in the north branch of the South-central drainage (Wetland 8).



Photo Point W8 WET DS – Photo looking southeast (downstream/off-site) of and wetland habitat in the north branch of the South-central drainage (Wetland 8) flowing toward Wetland 9.



Photo Point WP 9 UPL – Photo looking northwest (upstream/off-site) from stock pond berm of riparian and wetland habitat in the north branch of the South-central drainage (Wetland 9). Wetland 9 is connected to Wetland 8.



Photo Point WP 9 UPL – Photo looking southeast (downstream/on-site) from stock pond berm of riparian and wetland habitat in the north branch of the South-central drainage (Wetland 9) near Judge Orr Road.



Photo Point WP 9 WET – Photo looking west of wetland habitat where the north and south branches of the South-central drainage combine (Wetlands 8 and 9) adjacent to Judge Orr Road.



Photo Point WP 9 WET – Photo looking east of wetland habitat where the north and south branches of the South-central drainage combine (Wetlands 8 and 9) adjacent to Judge Orr Road.

Appendix C

USFWS IPaC Trust Resource Report

IPaC resource list

This report is an automatically generated list of species and other resources such as critical habitat (collectively referred to as *trust resources*) under the U.S. Fish and Wildlife Service's (USFWS) jurisdiction that are known or expected to be on or near the project area referenced below. The list may also include trust resources that occur outside of the project area, but that could potentially be directly or indirectly affected by activities in the project area. However, determining the likelihood and extent of effects a project may have on trust resources typically requires gathering additional site-specific (e.g., vegetation/species surveys) and project-specific (e.g., magnitude and timing of proposed activities) information.

Below is a summary of the project information you provided and contact information for the USFWS office(s) with jurisdiction in the defined project area. Please read the introduction to each section that follows (Endangered Species, Migratory Birds, USFWS Facilities, and NWI Wetlands) for additional information applicable to the trust resources addressed in that section.

Location

El Paso County, Colorado



Local office

Colorado Ecological Services Field Office

(303) 236-4773
(303) 236-4005

MAILING ADDRESS Denver Federal Center P.O. Box 25486 Denver, CO 80225-0486 PHYSICAL ADDRESS 134 Union Boulevard, Suite 670 Lakewood, CO 80228-1807

TFORCONSULTATION

Endangered species

This resource list is for informational purposes only and does not constitute an analysis of project level impacts.

The primary information used to generate this list is the known or expected range of each species. Additional areas of influence (AOI) for species are also considered. An AOI includes areas outside of the species range if the species could be indirectly affected by activities in that area (e.g., placing a dam upstream of a fish population even if that fish does not occur at the dam site, may indirectly impact the species by reducing or eliminating water flow downstream). Because species can move, and site conditions can change, the species on this list are not guaranteed to be found on or near the project area. To fully determine any potential effects to species, additional site-specific and projectspecific information is often required.

Section 7 of the Endangered Species Act **requires** Federal agencies to "request of the Secretary information whether any species which is listed or proposed to be listed may be present in the area of such proposed action" for any project that is conducted, permitted, funded, or licensed by any Federal agency. A letter from the local office and a species list which fulfills this requirement can **only** be obtained by requesting an official species list from either the Regulatory Review section in IPaC (see directions below) or from the local field office directly.

For project evaluations that require USFWS concurrence/review, please return to the IPaC website and request an official species list by doing the following:

- 1. Draw the project location and click CONTINUE.
- 2. Click DEFINE PROJECT.
- 3. Log in (if directed to do so).
- 4. Provide a name and description for your project.
- 5. Click REQUEST SPECIES LIST.

Listed species¹ and their critical habitats are managed by the <u>Ecological Services Program</u> of the U.S. Fish and Wildlife Service (USFWS) and the fisheries division of the National Oceanic and Atmospheric Administration (NOAA Fisheries²).

Species and critical habitats under the sole responsibility of NOAA Fisheries are **not** shown on this list. Please contact <u>NOAA Fisheries</u> for <u>species under their jurisdiction</u>.

- 1. Species listed under the <u>Endangered Species Act</u> are threatened or endangered; IPaC also shows species that are candidates, or proposed, for listing. See the <u>listing status page</u> for more information. IPaC only shows species that are regulated by USFWS (see FAQ).
- 2. <u>NOAA Fisheries</u>, also known as the National Marine Fisheries Service (NMFS), is an office of the National Oceanic and Atmospheric Administration within the Department of Commerce.

The following species are potentially affected by activities in this location:

Mammals

NAME	STATUS
 Gray Wolf Canis lupus This species only needs to be considered if the following condition applies: Lone, dispersing gray wolves may be present throughout the state of Colorado. If your activity includes a predator management program, please consider this species in your environmental review. There is final critical habitat for this species. https://ecos.fws.gov/ecp/species/4488 	Endangered
Preble's Meadow Jumping Mouse Zapus hudsonius preblei Wherever found There is final critical habitat for this species. Your location does not overlap the critical habitat. https://ecos.fws.gov/ecp/species/4090	Threatened
Birds NAME	STATUS
Eastern Black Rail Laterallus jamaicensis ssp. jamaicensis Wherever found No critical habitat has been designated for this species. https://ecos.fws.gov/ecp/species/10477	Threatened
 Piping Plover Charadrius melodus This species only needs to be considered if the following condition applies: Project includes water-related activities and/or use in the N. Platte, S. Platte, and Laramie River Basins which may affect listed species in Nebraska. There is final critical habitat for this species. Your location does not overlap the critical habitat. https://ecos.fws.gov/ecp/species/6039 	Threatened
Fishes	CTATUC
NAME	STATUS
Greenback Cutthroat Trout Oncorhynchus clarkii stomias	Threatened

Wherever found

No critical habitat has been designated for this species. <u>https://ecos.fws.gov/ecp/species/2775</u>

Pallid Sturgeon Scaphirhynchus albus

Endangered

Wherever found

This species only needs to be considered if the following condition applies:

 Project includes water-related activities and/or use in the N. Platte, S. Platte, and Laramie River Basins which may affect listed species in Nebraska.

No critical habitat has been designated for this species. <u>https://ecos.fws.gov/ecp/species/7162</u>

Insects

NAME	STATUS
Monarch Butterfly Danaus plexippus Wherever found No critical habitat has been designated for this species. <u>https://ecos.fws.gov/ecp/species/9743</u>	Candidate
Flowering Plants	STATUS
Ute Ladies'-tresses Spiranthes diluvialis Wherever found No critical habitat has been designated for this species. <u>https://ecos.fws.gov/ecp/species/2159</u>	Threatened

Critical habitats

Potential effects to critical habitat(s) in this location must be analyzed along with the endangered species themselves.

There are no critical habitats at this location.

You are still required to determine if your project(s) may have effects on all above listed species.

Migratory birds

Certain birds are protected under the Migratory Bird Treaty Act^{1} and the Bald and Golden Eagle Protection Act^{2} .

Any person or organization who plans or conducts activities that may result in impacts to migratory birds, eagles, and their habitats should follow appropriate regulations and consider implementing appropriate conservation measures, as described <u>below</u>.

1. The Migratory Birds Treaty Act of 1918.

https://ecos.fws.gov/ecp/species/6038

2. The <u>Bald and Golden Eagle Protection Act</u> of 1940.

Additional information can be found using the following links:

- Birds of Conservation Concern https://www.fws.gov/program/migratory-birds/species
- Measures for avoiding and minimizing impacts to birds <u>https://www.fws.gov/library/collections/avoiding-and-minimizing-incidental-take-migratory-birds</u>
- Nationwide conservation measures for birds <u>https://www.fws.gov/sites/default/files/documents/nationwide-standard-conservation-measures.pdf</u>

The birds listed below are birds of particular concern either because they occur on the <u>USFWS</u> <u>Birds of Conservation Concern</u> (BCC) list or warrant special attention in your project location. To learn more about the levels of concern for birds on your list and how this list is generated, see the FAQ <u>below</u>. This is not a list of every bird you may find in this location, nor a guarantee that every bird on this list will be found in your project area. To see exact locations of where birders and the general public have sighted birds in and around your project area, visit the <u>E-bird data mapping tool</u> (Tip: enter your location, desired date range and a species on your list). For projects that occur off the Atlantic Coast, additional maps and models detailing the relative occurrence and abundance of bird species on your list are available. Links to additional information about Atlantic Coast birds, and other important information about your migratory bird list, including how to properly interpret and use your migratory bird report, can be found <u>below</u>.

For guidance on when to schedule activities or implement avoidance and minimization measures to reduce impacts to migratory birds on your list, click on the PROBABILITY OF PRESENCE SUMMARY at the top of your list to see when these birds are most likely to be present and breeding in your project area.

NAME	BREEDING SEASON
Bald Eagle Haliaeetus leucocephalus This is not a Bird of Conservation Concern (BCC) in this area, but warrants attention because of the Eagle Act or for potential susceptibilities in offshore areas from certain types of development or activities.	Breeds Oct 15 to Jul 31
Ferruginous Hawk Buteo regalis This is a Bird of Conservation Concern (BCC) only in particular Bird Conservation Regions (BCRs) in the continental USA	Breeds Mar 15 to Aug 15

Golden Eagle Aquila chrysaetos This is not a Bird of Conservation Concern (BCC) in this area, but warrants attention because of the Eagle Act or for potential susceptibilities in offshore areas from certain types of development or activities. <u>https://ecos.fws.gov/ecp/species/1680</u>

Lesser Yellowlegs Tringa flavipes This is a Bird of Conservation Concern (BCC) throughout its range in the continental USA and Alaska. <u>https://ecos.fws.gov/ecp/species/9679</u>

Pinyon Jay Gymnorhinus cyanocephalus This is a Bird of Conservation Concern (BCC) throughout its range in the continental USA and Alaska. <u>https://ecos.fws.gov/ecp/species/9420</u> Breeds Feb 15 to Jul 15

Probability of Presence Summary

The graphs below provide our best understanding of when birds of concern are most likely to be present in your project area. This information can be used to tailor and schedule your project activities to avoid or minimize impacts to birds. Please make sure you read and understand the FAQ "Proper Interpretation and Use of Your Migratory Bird Report" before using or attempting to interpret this report.

Probability of Presence (

Each green bar represents the bird's relative probability of presence in the 10km grid cell(s) your project overlaps during a particular week of the year. (A year is represented as 12 4-week months.) A taller bar indicates a higher probability of species presence. The survey effort (see below) can be used to establish a level of confidence in the presence score. One can have higher confidence in the presence score if the corresponding survey effort is also high.

How is the probability of presence score calculated? The calculation is done in three steps:

- 1. The probability of presence for each week is calculated as the number of survey events in the week where the species was detected divided by the total number of survey events for that week. For example, if in week 12 there were 20 survey events and the Spotted Towhee was found in 5 of them, the probability of presence of the Spotted Towhee in week 12 is 0.25.
- 2. To properly present the pattern of presence across the year, the relative probability of presence is calculated. This is the probability of presence divided by the maximum probability of presence across all weeks. For example, imagine the probability of presence in week 20 for the Spotted Towhee is 0.05, and that the probability of presence at week 12 (0.25) is the maximum of any week of the year. The relative probability of presence on week 12 is 0.25/0.25 = 1; at week 20 it is 0.05/0.25 = 0.2.
- 3. The relative probability of presence calculated in the previous step undergoes a statistical conversion so that all possible values fall between 0 and 10, inclusive. This is the probability of presence score.

Breeds elsewhere

To see a bar's probability of presence score, simply hover your mouse cursor over the bar.

Breeding Season (=)

Yellow bars denote a very liberal estimate of the time-frame inside which the bird breeds across its entire range. If there are no yellow bars shown for a bird, it does not breed in your project area.

Survey Effort (l)

Vertical black lines superimposed on probability of presence bars indicate the number of surveys performed for that species in the 10km grid cell(s) your project area overlaps. The number of surveys is expressed as a range, for example, 33 to 64 surveys.

To see a bar's survey effort range, simply hover your mouse cursor over the bar.

No Data (–)

A week is marked as having no data if there were no survey events for that week.

Survey Timeframe

Surveys from only the last 10 years are used in order to ensure delivery of currently relevant information. The exception to this is areas off the Atlantic coast, where bird returns are based on all years of available data, since data in these areas is currently much more sparse.

				■ prob	ability of	presen	ce 📕 br	eeding se	eason	survey	effort -	– no data
SPECIES	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
Bald Eagle Non-BCC Vulnerable	+1+1	++1+	++∎+	1+++	H++++	++++	THI	++++	++++	++++	+11+	++1+
Ferruginous Hawk BCC - BCR	++	+ -	+++++	++(+	++++	++++	+++#	++++	++++	++++	++++	++++
Golden Eagle Non-BCC Vulnerable	+1+1		Ŷť	++1+	++++	++++	++I+	++++	++++	++++	+++‡	++++
Lesser Yellowlegs BCC Rangewide (CON)	++++	++++	++++	+∎∎∎	++++	++++	++++	++++	++++	++++	++++	++++
Pinyon Jay BCC Rangewide (CON)	++++	++++	++++	++++	++++	++++	++++	++++	++++	+++	++++	++++

Tell me more about conservation measures I can implement to avoid or minimize impacts to migratory birds.

<u>Nationwide Conservation Measures</u> describes measures that can help avoid and minimize impacts to all birds at any location year round. Implementation of these measures is particularly important when birds are most likely to occur in the project area. When birds may be breeding in the area, identifying the locations of any active nests and avoiding their destruction is a very helpful impact minimization measure. To see when birds are most likely to occur and be breeding in your project area, view the Probability of Presence Summary. <u>Additional measures</u> or <u>permits</u> may be advisable depending on the type of activity you are conducting and the type of infrastructure or bird species present on your project site.

The Migratory Bird Resource List is comprised of USFWS <u>Birds of Conservation Concern (BCC)</u> and other species that may warrant special attention in your project location.

The migratory bird list generated for your project is derived from data provided by the <u>Avian Knowledge Network</u> (<u>AKN</u>). The AKN data is based on a growing collection of <u>survey</u>, <u>banding</u>, <u>and citizen science datasets</u> and is queried and filtered to return a list of those birds reported as occurring in the 10km grid cell(s) which your project intersects, and that have been identified as warranting special attention because they are a BCC species in that area, an eagle (<u>Eagle Act</u> requirements may apply), or a species that has a particular vulnerability to offshore activities or development.

Again, the Migratory Bird Resource list includes only a subset of birds that may occur in your project area. It is not representative of all birds that may occur in your project area. To get a list of all birds potentially present in your project area, please visit the <u>Rapid Avian Information Locator (RAIL) Tool</u>.

What does IPaC use to generate the probability of presence graphs for the migratory birds potentially occurring in my specified location?

The probability of presence graphs associated with your migratory bird list are based on data provided by the <u>Avian</u> <u>Knowledge Network (AKN)</u>. This data is derived from a growing collection of <u>survey</u>, <u>banding</u>, <u>and citizen science</u> <u>datasets</u>.

Probability of presence data is continuously being updated as new and better information becomes available. To learn more about how the probability of presence graphs are produced and how to interpret them, go the Probability of Presence Summary and then click on the "Tell me about these graphs" link.

How do I know if a bird is breeding, wintering or migrating in my area?

To see what part of a particular bird's range your project area falls within (i.e. breeding, wintering, migrating or yearround), you may query your location using the <u>RAIL Tool</u> and look at the range maps provided for birds in your area at the bottom of the profiles provided for each bird in your results. If a bird on your migratory bird species list has a breeding season associated with it, if that bird does occur in your project area, there may be nests present at some point within the timeframe specified. If "Breeds elsewhere" is indicated, then the bird likely does not breed in your project area.

What are the levels of concern for migratory birds?

Migratory birds delivered through IPaC fall into the following distinct categories of concern:

- 1. "BCC Rangewide" birds are <u>Birds of Conservation Concern</u> (BCC) that are of concern throughout their range anywhere within the USA (including Hawaii, the Pacific Islands, Puerto Rico, and the Virgin Islands);
- 2. "BCC BCR" birds are BCCs that are of concern only in particular Bird Conservation Regions (BCRs) in the continental USA; and
- 3. "Non-BCC Vulnerable" birds are not BCC species in your project area, but appear on your list either because of the <u>Eagle Act</u> requirements (for eagles) or (for non-eagles) potential susceptibilities in offshore areas from certain types of development or activities (e.g. offshore energy development or longline fishing).

Although it is important to try to avoid and minimize impacts to all birds, efforts should be made, in particular, to avoid and minimize impacts to the birds on this list, especially eagles and BCC species of rangewide concern. For more information on conservation measures you can implement to help avoid and minimize migratory bird impacts and requirements for eagles, please see the FAQs for these topics.

Details about birds that are potentially affected by offshore projects

For additional details about the relative occurrence and abundance of both individual bird species and groups of bird species within your project area off the Atlantic Coast, please visit the <u>Northeast Ocean Data Portal</u>. The Portal also offers data and information about other taxa besides birds that may be helpful to you in your project review. Alternately, you may download the bird model results files underlying the portal maps through the <u>NOAA NCCOS</u> <u>Integrative Statistical Modeling and Predictive Mapping of Marine Bird Distributions and Abundance on the Atlantic Outer Continental Shelf</u> project webpage.

Bird tracking data can also provide additional details about occurrence and habitat use throughout the year, including migration. Models relying on survey data may not include this information. For additional information on marine bird tracking data, see the <u>Diving Bird Study</u> and the <u>nanotag studies</u> or contact <u>Caleb Spiegel</u> or <u>Pam Loring</u>.

What if I have eagles on my list?

If your project has the potential to disturb or kill eagles, you may need to <u>obtain a permit</u> to avoid violating the Eagle Act should such impacts occur.

Proper Interpretation and Use of Your Migratory Bird Report

The migratory bird list generated is not a list of all birds in your project area, only a subset of birds of priority concern. To learn more about how your list is generated, and see options for identifying what other birds may be in your project area, please see the FAQ "What does IPaC use to generate the migratory birds potentially occurring in my specified location". Please be aware this report provides the "probability of presence" of birds within the 10 km grid cell(s) that overlap your project; not your exact project footprint. On the graphs provided, please also look carefully at the survey effort (indicated by the black vertical bar) and for the existence of the "no data" indicator (a red horizontal bar). A high survey effort is the key component. If the survey effort is high, then the probability of presence score can be viewed as more dependable. In contrast, a low survey effort bar or no data bar means a lack of data and, therefore, a lack of certainty about presence of the species. This list is not perfect; it is simply a starting point for identifying what birds of concern have the potential to be in your project area, when they might be there, and if they might be breeding (which means nests might be present). The list helps you know what to look for to confirm presence, and helps guide you in knowing when to implement conservation measures to avoid or minimize potential impacts from your project activities, should presence be confirmed. To learn more about conservation measures, visit the FAQ "Tell me about conservation measures I can implement to avoid or minimize impacts to migratory birds" at the bottom of your migratory bird trust resources page.

Facilities

National Wildlife Refuge lands

Any activity proposed on lands managed by the <u>National Wildlife Refuge</u> system must undergo a 'Compatibility Determination' conducted by the Refuge. Please contact the individual Refuges to discuss any questions or concerns.

There are no refuge lands at this location.

Fish hatcheries

There are no fish hatcheries at this location.

Wetlands in the National Wetlands Inventory (NWI)

Impacts to <u>NWI wetlands</u> and other aquatic habitats may be subject to regulation under Section 404 of the Clean Water Act, or other State/Federal statutes.

For more information please contact the Regulatory Program of the local <u>U.S. Army Corps of</u> <u>Engineers District</u>.

Wetland information is not available at this time

This can happen when the National Wetlands Inventory (NWI) map service is unavailable, or for very large projects that intersect many wetland areas. Try again, or visit the <u>NWI map</u> to view wetlands at this location.

Data limitations

The Service's objective of mapping wetlands and deepwater habitats is to produce reconnaissance level information on the location, type and size of these resources. The maps are prepared from the analysis of high altitude imagery. Wetlands are identified based on vegetation, visible hydrology and geography. A margin of error is inherent in the use of imagery; thus, detailed on-the-ground inspection of any particular site may result in revision of the wetland boundaries or classification established through image analysis.

The accuracy of image interpretation depends on the quality of the imagery, the experience of the image analysts, the amount and quality of the collateral data and the amount of ground truth verification work conducted. Metadata should be consulted to determine the date of the source imagery used and any mapping problems.

Wetlands or other mapped features may have changed since the date of the imagery or field work. There may be occasional differences in polygon boundaries or classifications between the information depicted on the map and the actual conditions on site.

Data exclusions

Certain wetland habitats are excluded from the National mapping program because of the limitations of aerial imagery as the primary data source used to detect wetlands. These habitats include seagrasses or submerged aquatic vegetation that are found in the intertidal and subtidal zones of estuaries and nearshore coastal waters. Some deepwater reef communities (coral or tuberficid worm reefs) have also been excluded from the inventory. These habitats, because of their depth, go undetected by aerial imagery.

Data precautions

Federal, state, and local regulatory agencies with jurisdiction over wetlands may define and describe wetlands in a different manner than that used in this inventory. There is no attempt, in either the design or products of this inventory, to define the limits of proprietary jurisdiction of any Federal, state, or local government or to establish the geographical scope of the regulatory programs of government agencies. Persons intending to engage in activities involving modifications within or adjacent to wetland areas should seek the advice of appropriate Federal, state, or local agencies concerning specified agency regulatory programs and proprietary jurisdictions that may affect such activities.

CONSUL

OR

Appendix D

USACE Wetland Determination Data Forms

Project/Site: Davis	City/County: EL Page Sampling Date: 5/22/23
Applicant/Owner: Davis Investigator(s): Jen Davzyavdis+Grvant Grvmee:	State: <u>CO</u> Sampling Point: <u>W4-WE</u> Section, Township, Range: <u>S33434</u> , T125, R64W
Subregion (LRR): G	disturbed? Are "Normal Circumstances" present? Yes No
Hydrophytic Vegetation Present? Yes No Hydric Soil Present? Yes No Wetland Hydrology Present? Yes No Remarks: No No	Is the Sampled Area within a Wetland? Yes No

VEGETATION - Use scientific names of plants.

Tree Stratum (Plot size:) 1		Dominant Species?	Status	Dominance Test worksheet: Number of Dominant Species That Are OBL, FACW, or FAC (excluding FAC-): Total Number of Dominant Species Across All Strata: (B)
4	20	= Total Cov	er OBL	Percent of Dominant Species 83.33 (A/B)
2				Prevalence Index worksheet: Total % Cover of:Muttiply by:
3				Total % Cover of Multiply by: OBL species x 1 =
4				FACW species x 2 =
5				FAC species x 3 =
Herb Stratum (Plot size:) 1balticus 2aurex nebrascensis 3aurex praegracilis 4aurex praegracilis 5aurense 6ypha angustitalia 7 8	25	= Total Cov		FAC species x 3 = FACU species x 4 = UPL species x 5 = Column Totals: (A) (B) Prevalence Index = B/A = Hydrophytic Vegetation Indicators: 1 - Rapid Test for Hydrophytic Vegetation 2 - Dominance Test is >50% 3 - Prevalence Index is $\leq 3.0^1$ 4 - Morphological Adaptations ¹ (Provide supporting data in Remarks or on a separate sheet)
10	90	= Total Cov	er	Problematic Hydrophytic Vegetation ¹ (Explain) ¹ Indicators of hydric soil and wetland hydrology must be present, unless disturbed or problematic.
2	100	= Total Cov	er	Hydrophytic Vegetation Present? Yes <u>No</u> No
Remarks:				

Sampling Point: WY-WET

Profile Desci	ription: (Descri	be to the dep	th needed to docum	nent the indica	tor or confirm	the absence	of indicators.)
Depth	Matri		Redo	Features			
(inches)	Color (moist)	%	Color (moist)	<u>%</u> Typ	e ¹ Loc ²	Texture	Remarks
0-2	N/F					OM	Organic Matter
2-10	10 YR 2.51	N a				Sand	0
0-105	104R 51	1 75	IDVR512	25 0	M	Sitty Cla	1/
10-10	10/12/1		10/1-0/10	2		still and	9
0. 17			······································				······································
	<u> </u>						
			Reduced Matrix, CS		ated Sand Gr	and the Report of Contract of	cation: PL=Pore Lining, M=Matrix.
		licable to all	LRRs, unless other	wise noted.)		Indicators	for Problematic Hydric Soils ³ :
Histosol (eleyed Matrix (S	4)	1 cm M	Muck (A9) (LRR I, J)
	ipedon (A2)			edox (S5)			Prairie Redox (A16) (LRR F, G, H)
Black His				Matrix (S6)			Surface (S7) (LRR G)
	n Sulfide (A4)			Aucky Mineral (F			Plains Depressions (F16)
	Layers (A5) (LR			Bleyed Matrix (F	2)		RR H outside of MLRA 72 & 73)
요즘 그는 아이들은 것은 것을 했다.	ck (A9) (LRR F, Below Dark Sur			Matrix (F3)	×	A COLOR	ed Vertic (F18)
	rk Surface (A12)			ark Surface (F6 Dark Surface (the second s	arent Material (TF2) Shallow Dark Surface (TF12)
	ucky Mineral (S1			epressions (F8		Contraction of the Contraction	(Explain in Remarks)
	lucky Peat or Pe		S AGE (0.3265) 325	ins Depressions			of hydrophytic vegetation and
	cky Peat or Peat			RA 72 & 73 of L			d hydrology must be present,
		A	1.00				disturbed or problematic.
Restrictive L	ayer (if present):				1.	
Type:	28 2. 48 3						
Depth (incl	hes):					Hydric Soil	Present? Yes V No
Remarks:						1.	
IYDROLOG	GY						
Wetland Hyd	rology Indicato	rs:					
a series of the	Constraint and the second statement of the		: check all that apply)		Seconda	ary Indicators (minimum of two required)
. /	Water (A1)		Salt Crust			Line Contraction	face Soil Cracks (B6)
	er Table (A2)			ertebrates (B13	1		rsely Vegetated Concave Surface (B8)
V Saturation				Sulfide Odor (C1			
Water Ma				Water Table (Comments States	inage Patterns (B10)
	t Deposits (B2)				State and the o		dized Rhizospheres on Living Roots (C3)
Drift Depo			STATE STREET	hizospheres on	Living Roots ((here tilled)
Name of the second second second second	or Crust (B4)		(where n				yfish Burrows (C8)
Iron Depo			the second second second second	f Reduced Iron Surface (C7)	(04)		aration Visible on Aerial Imagery (C9)
the share and the	n Visible on Aeri	al Imagon (P	the second s	ain in Remarks		N C	morphic Position (D2)
STATE AND A STATE) Other (Exp	ain in Remarks			C-Neutral Test (D5)
Field Observa	ained Leaves (B	")				Fros	st-Heave Hummocks (D7) (LRR F)
		1		hes): 0-1 m	6		
Surface Wate			No Depth (inc		0		
Water Table F		. /	No Depth (inc	0			/
Saturation Pre		Yes V	No Depth (inc	hes): 🥖	Wetla	and Hydrology	y Present? Yes V No
(includes capi Describe Reco		am gauge, mo	nitoring well, aerial p	hotos, previous	inspections) i	f available:	
			the state of the s			a a randolo.	
Pomorke			11	1.0			
Remarks:	Broad	SWO	th of fl	odolar	1		
	01000	00000		A acce			

Project/Site: Davis	City/County: El Pag	50 5	Sampling Date:	122/23
Applicant/Owner: Davis		State: <u>CD</u> s	Sampling Point: 🚺	4-UPL
Investigator(s): Jon Davzvardis+Grant Gromee	Section, Township, Range:	: 533+34, T12:	S,RGYW	
Landform (hillslope, terrace, etc.):	Local relief (concave, conv		Slope	(%):0-3
		ong: -104,543=		WGS84
Soil Map Unit Name: 19- Colvmbne gravelly same	ly loam, 0-3% slop	NWI classificat	tion: N/A	
Are climatic / hydrologic conditions on the site typical for this time of ye	ar? Yes No	(If no, explain in Rer	marks.)	
Are Vegetation, Soil, or Hydrology significantly	disturbed? Are "Nor	mal Circumstances" pre	esent? Yes	No
Are Vegetation, Soil, or Hydrology naturally pro	oblematic? (If neede	d, explain any answers	in Remarks.)	
SUMMARY OF FINDINGS – Attach site map showing	sampling point loca	ations, transects, i	important feat	ures, etc.

Hydrophytic Vegetation Present? Hydric Soil Present? Wetland Hydrology Present?	Yes Yes Yes	Is the Sampled Area within a Wetland?	Yes	No
Remarks:		1		
· •				

VEGETATION - Use scientific names of plants.

<u>Tree Stratum</u> (Plot size:) 1)		Species?	Status	Dominance Test worksheet: Number of Dominant Species That Are OBL, FACW, or FAC
2				(excluding FAC-): (A)
3				Total Number of Dominant Species Across All Strata: (B)
Sapling/Shrub Stratum (Plot size:) 1)		= Total Cov		Percent of Dominant Species That Are OBL, FACW, or FAC: (A/
2				Prevalence Index worksheet:
3				Total % Cover of: Multiply by:
4				OBL species x 1 =
5				FACW species x 2 =
		= Total Cov	er	FAC species x 3 =
Herb Stratum (Plot size:)				FACU species x 4 =
1. Bouteloua gracitic	- 40	Y	UPL	UPL species x 5 =
2				Column Totals: (A) (B
3				Prevalence Index = B/A =
4				
5				Hydrophytic Vegetation Indicators: 1 - Rapid Test for Hydrophytic Vegetation
6				2 - Dominance Test is >50%
7				$\frac{2}{3} - \text{Prevalence Index is } \le 3.0^{1}$
8				(V-50)
9				4 - Morphological Adaptations ¹ (Provide supportin data in Remarks or on a separate sheet)
10				Problematic Hydrophytic Vegetation ¹ (Explain)
Woody Vine Stratum (Plot size:) 1)		= Total Cov		¹ Indicators of hydric soil and wetland hydrology must be present, unless disturbed or problematic.
2		= Total Cov		Hydrophytic Vegetation
% Bare Ground in Herb Stratum			8 	Present? Yes No V
Remarks:				
JS Army Corps of Engineers				Great Plains – Version 2.

Sampling Point: My-UPL

Profile Description: (Describe to the depth		confirm the absence of indicators.)
Depth <u>Matrix</u>	Redox Features	
(inches) Color (moist) %	Color (moist) % Type ¹ I	Loc ² Texture Remarks
0-10 10483/ 100 -		ClayLoam Dry
16-18 10 YR3/ 6 100		Sandy C/ Lm
		······································
¹ Type: C=Concentration, D=Depletion, RM=F		
Hydric Soil Indicators: (Applicable to all L		Indicators for Problematic Hydric Soils ³ :
Histosol (A1)	Sandy Gleyed Matrix (S4)	1 cm Muck (A9) (LRR I, J)
Histic Epipedon (A2)	Sandy Redox (S5)	Coast Prairie Redox (A16) (LRR F, G, H)
Black Histic (A3)	Stripped Matrix (S6)	Dark Surface (S7) (LRR G)
Hydrogen Sulfide (A4) Stratified Layers (A5) (LRR F)	Loamy Mucky Mineral (F1) Loamy Gleyed Matrix (F2)	High Plains Depressions (F16)
1 cm Muck (A9) (LRR F, G, H)	Depleted Matrix (F2)	(LRR H outside of MLRA 72 & 73) Reduced Vertic (F18)
Depleted Below Dark Surface (A11)	Redox Dark Surface (F6)	Red Parent Material (TF2)
Thick Dark Surface (A12)	Depleted Dark Surface (F7)	Very Shallow Dark Surface (TF12)
Sandy Mucky Mineral (S1)	Redox Depressions (F8)	Other (Explain in Remarks)
2.5 cm Mucky Peat or Peat (S2) (LRR G,	H) High Plains Depressions (F16)	
5 cm Mucky Peat or Peat (S3) (LRR F)	(MLRA 72 & 73 of LRR H)	wetland hydrology must be present,
Destriction I and the second		unless disturbed or problematic.
Restrictive Layer (if present):		
Type:		1/1
Depth (inches):		Hydric Soil Present? Yes No
Remarks:		
HYDROLOGY		
Wetland Hydrology Indicators:		
Primary Indicators (minimum of one required;	check all that apply)	Secondary Indicators (minimum of two required)
Surface Water (A1)	Salt Crust (B11)	Surface Soil Cracks (B6)
High Water Table (A2)	Aquatic Invertebrates (B13)	Sparsely Vegetated Concave Surface (B8)
Saturation (A3)	Hydrogen Sulfide Odor (C1)	Drainage Patterns (B10)
Water Marks (B1)	Dry-Season Water Table (C2)	Oxidized Rhizospheres on Living Roots (C3)
Sediment Deposits (B2)	Oxidized Rhizospheres on Living	Roots (C3) (where tilled)
Drift Deposits (B3)	(where not tilled)	Crayfish Burrows (C8)
Algal Mat or Crust (B4)	Presence of Reduced Iron (C4)	Saturation Visible on Aerial Imagery (C9)
Iron Deposits (B5)	Thin Muck Surface (C7)	Geomorphic Position (D2)
Inundation Visible on Aerial Imagery (B7)	Other (Explain in Remarks)	FAC-Neutral Test (D5)
Water-Stained Leaves (B9)	/	Frost-Heave Hummocks (D7) (LRR F)
Field Observations:		
	Depth (inches):	7
Water Table Present? Yes No	Depth (inches):	
	Depth (inches):	Wetland Hydrology Present? Yes No
(includes capillary fringe) Describe Recorded Data (stream gauge, moni	toring well aerial photos, previous insper	tions) if available:
	toring won, donal protos, previous lisped	diono, il avaliable.
Remarks:		

Project/Site: Davis	City/County: El Pago Sampling	Date: 5/22/23
Applicant/Owner: Davis	State: <u>CD</u> Sampling	Point: W5-WET
Investigator(s): Jon Davzyavais+Grvant Grvmee	Section, Township, Range: 533434, T125, RG	YW .
Landform (hillslope, terrace, etc.): <u>hillslope</u>	Local relief (concave, convex, none): Concave	
Subregion (LRR): Lat:	8-961883 N Long: -104,543390°	Watum: WG-S 84
Soil Map Unit Name: 19-Columbing gravely sandy	learn, 0-3% slepes_ NWI classification: F	EM JA
Are climatic / hydrologic conditions on the site typical for this time of ye	ar? Yes No (If no, explain in Remarks.)	1
Are Vegetation, Soil, or Hydrology significantly	disturbed? Are "Normal Circumstances" present? Y	es No
Are Vegetation, Soil, or Hydrology naturally pro	blematic? (If needed, explain any answers in Reman	ks.)
SUMMARY OF FINDINGS – Attach site map showing	sampling point locations, transects, importa	int features, etc.

Hydrophytic Vegetation Present? Hydric Soil Present? Wetland Hydrology Present?	Yes No Yes No Yes No	Is the Sampled Area within a Wetland?	Yes No
Remarks:			2,
		- C	
10 et			

VEGETATION - Use scientific names of plants.

Tree Stratum (Plot size:) 1)		Dominant Indicator Species? Status	Dominance Test worksheet: Number of Dominant Species That Are OBL, FACW, or FAC
2			(excluding FAC-): (A)
3			Total Number of Dominant
4			Species Across All Strata:
Sapling/Shrub Stratum (Plot size:) 1		= Total Cover	Percent of Dominant Species That Are OBL, FACW, or FAC:
2			Prevalence Index worksheet:
3			Total % Cover of: Multiply by:
4			OBL species x 1 =
5			FACW species x 2 =
		= Total Cover	FAC species x 3 =
Herb Stratum (Plot size:)			FACU species x 4 =
1. Juncus balticus.	40	Y OBL	UPL species x 5 =
2. Carex ne braiscensis	25	Y OBL	Column Totals: (A) (B)
3. Tris Missouriensis	P	N OBL	
4. Cirbium anvense	0	Y FACU	Prevalence Index = B/A =
5. Covex proparaellis	25	V OBL	Hydrophytic Vegetation Indicators:
6		/	1 - Rapid Test for Hydrophytic Vegetation
7			✓ 2 - Dominance Test is >50%
8			3 - Prevalence Index is ≤3.0 ¹
9			4 - Morphological Adaptations ¹ (Provide supporting data in Remarks or on a separate sheet)
10.			Problematic Hydrophytic Vegetation ¹ (Explain)
	100	= Total Cover	
Woody Vine Stratum (Plot size:)			¹ Indicators of hydric soil and wetland hydrology must be present, unless disturbed or problematic.
1			be present, unless disturbed or problematic.
2	10.0		Hydrophytic
% Bare Ground in Herb Stratum	100	= Total Cover	Vegetation Present? Yes No No
Remarks:			

Sampling Point:	W3-	WE
-----------------	-----	----

Profile Description: (Describe to the depth need	eded to document the indicator	or confirm	n the absence of indicators.)
Depth <u>Matrix</u>	Redox Features		
	olor (moist) % Type ¹	Loc ²	Texture Remarks
0-9 10YR3/1 100			Silty Clay Loan
8-18 104R 2/1 100			Silty Clay
			terre and the second se
¹ Type: C=Concentration, D=Depletion, RM=Redu		d Sand G	
Hydric Soil Indicators: (Applicable to all LRRs	and a second		Indicators for Problematic Hydric Soils ³ :
Histosol (A1) Histic Epipedon (A2)	Sandy Gleyed Matrix (S4)		1 cm Muck (A9) (LRR I, J)
Black Histic (A3)	Sandy Redox (S5) Stripped Matrix (S6)		Coast Prairie Redox (A16) (LRR F, G, H)
Hydrogen Sulfide (A4)	Loamy Mucky Mineral (F1)		Dark Surface (S7) (LRR G) High Plains Depressions (F16)
Stratified Layers (A5) (LRR F)	Loamy Gleyed Matrix (F2)		(LRR H outside of MLRA 72 & 73)
1 cm Muck (A9) (LRR F, G, H)	Depleted Matrix (F3)		Reduced Vertic (F18)
Depleted Below Dark Surface (A11)	Redox Dark Surface (F6)		Red Parent Material (TF2)
Thick Dark Surface (A12)	Depleted Dark Surface (F7)		Very Shallow Dark Surface (TF12)
Sandy Mucky Mineral (S1)	Redox Depressions (F8)		Other (Explain in Remarks)
2.5 cm Mucky Peat or Peat (S2) (LRR G, H)	High Plains Depressions (F		³ Indicators of hydrophytic vegetation and
5 cm Mucky Peat or Peat (S3) (LRR F)	(MLRA 72 & 73 of LRR	H)	wetland hydrology must be present,
Restrictive Layer (if present):			unless disturbed or problematic.
Depth (inches):			
		_	Hydric Soil Present? Yes V No
Remarks:			
HYDROLOGY			
Wetland Hydrology Indicators:			
Primary Indicators (minimum of one required; chec	k all that apply)		Secondary Indicators (minimum of two required)
Surface Water (A1)	Salt Crust (B11)		
High Water Table (A2)	Aquatic Invertebrates (B13)		Surface Soil Cracks (B6)
Saturation (A3)	Hydrogen Sulfide Odor (C1)		Sparsely Vegetated Concave Surface (B8)
Water Marks (B1)	Dry-Season Water Table (C2)		Drainage Patterns (B10)
Sediment Deposits (B2)	 Oxidized Rhizospheres on Livi 	a Poote (Oxidized Rhizospheres on Living Roots (C3)
Drift Deposits (B3)	(where not tilled)	ig noois (C3) (where tilled) Crayfish Burrows (C8)
Algal Mat or Crust (B4)	Presence of Reduced Iron (C4		Saturation Visible on Aerial Imagery (C9)
Iron Deposits (B5)	Thin Muck Surface (C7)		Geomorphic Position (D2)
Inundation Visible on Aerial Imagery (B7)	Other (Explain in Remarks)		FAC-Neutral Test (D5)
Water-Stained Leaves (B9)			Frost-Heave Hummocks (D7) (LRR F)
Field Observations:			
Surface Water Present? Yes No V	Depth (inches):		
Water Table Present? Yes V	Depth (inches): 10 inches	5	2
Saturation Present? Yes V No	Depth (inches): 10 unches	Wotls	Ind Hydrology Present? Yes No
(includes capillary fringe)			
Describe Recorded Data (stream gauge, monitorin	g well, aerial photos, previous insp	ections), i	f available:
Remarks:) (1	1	
Saturated soil down	stream of low	vest	Steck pond
and a surround a surround of			

Project/Site:	_ City/County: <u>El Pago</u> Sampling Date: <u>5/22/23</u>
Applicant/Owner: Davis	State: CO Sampling Point: W5-UPL
Investigator(s): Jon Davzyavdist Givant Givmet	2 Section, Township, Range: <u>533434, T125, R64W</u>
Landform (hillslope, terrace, etc.):hllglope	_ Local relief (concave, convex, none): <u>CCNCave</u> Slope (%): <u>0-3</u>
Subregion (LRR): Lat:	38.96[983" Long: -104,543390 Watum: WGS 84
Soil Map Unit Name: 19- Columbre gravelly soud loa	M, 0-326 Slopes NWI classification: N/A
Are climatic / hydrologic conditions on the site typical for this time of	year? Yes No (If no, explain in Remarks.)
Are Vegetation, Soil, or Hydrology significant	tly disturbed? Are "Normal Circumstances" present? Yes No
Are Vegetation, Soil, or Hydrology naturally	problematic? (If needed, explain any answers in Remarks.)
SUMMARY OF FINDINGS - Attach site map showin	ng sampling point locations, transects, important features, etc.
Hydrophytic Vegetation Present? Yes No	is the Completion
Hydric Soil Present? Yes No	Is the Sampled Area within a Wetland? Yes No
Wetland Hydrology Present? Yes No	
Remarks:	

VEGETATION - Use scientific names of plants.

Tree Stratum (Plot size:)	Absolute % Cover	Dominant Indicator Species? Status	Dominance Test worksheet: Number of Dominant Species
1			That Are ORL EACIN or EAC
2			(excluding FAC-): (A)
3			Total Number of Dominant
4			Species Across All Strata: (B)
Sapling/Shrub Stratum (Plot size:)		= Total Cover	Percent of Dominant Species That Are OBL, FACW, or FAC: (A/B)
12			Prevalence Index worksheet:
2 3			Total % Cover of. Multiply by:
4			OBL species x 1 =
5			FACW species x 2 =
	-	= Total Cover	FAC species x 3 =
Herb Stratum (Plot size:)		- Total Cover	FACU species x 4 =
1. Bromus inermis	100	Y UPL	UPL species x 5 =
2			Column Totals: (A) (B)
3			Provide the Provid
4			Prevalence Index = B/A =
5			Hydrophytic Vegetation Indicators:
6			1 - Rapid Test for Hydrophytic Vegetation
7			2 - Dominance Test is >50%
8			3 - Prevalence Index is ≤3.0 ¹
9			4 - Morphological Adaptations ¹ (Provide supporting data in Remarks or on a separate sheet)
10			Problematic Hydrophytic Vegetation ¹ (Explain)
	100	= Total Cover	
Woody Vine Stratum (Plot size:) 1)			¹ Indicators of hydric soil and wetland hydrology must be present, unless disturbed or problematic.
2			Hydrophytic /
% Bare Ground in Herb Stratum	100	= Total Cover	Vegetation Present? Yes No
Remarks:		and the second	

Sampling Point:	W5-	UPL
		_

	ription: (Describe	to the depth n	eeded to docum	nent the	indicator o	or confirm	m the absence of indicators.)
Depth (inchor)	Matrix	0/		x Feature			
(inches)	Color (moist)	<u> % </u>	Color (moist)	%	_Type ¹	Loc ²	Texture Remarks
0-10	104K21						Clay Leam Dry
16-18	107R312	100	1-010-00-00-00-00-00-00-00-00-00-00-00-0		-	0	Sandy Clicam
							· · · · · · · · · · · · · · · · · · ·
						_	
¹ Type: C=Co	oncentration, D=Dep	pletion. RM=Rec	luced Matrix CS	=Covered	or Coater	Sand G	rains. ² Location: PL=Pore Lining, M=Mätrix.
Hydric Soil I	ndicators: (Applic	able to all LRR	s, unless other	wise not	ed.)		Indicators for Problematic Hydric Soils ³ :
Histosol				ileyed Ma	and the second second		1 cm Muck (A9) (LRR I, J)
Histic Ep	ipedon (A2)			edox (S5			Coast Prairie Redox (A16) (LRR F, G, H)
Black His				Matrix (S			Dark Surface (S7) (LRR G)
	n Sulfide (A4)		contraction of the second states of the second stat		eral (F1)	/	High Plains Depressions (F16)
	Layers (A5) (LRR			Heyed Ma	and the second		(LRR H outside of MLRA 72 & 73)
	ck (A9) (LRR F, G, Below Dark Surfac			Matrix (F			Reduced Vertic (F18)
	rk Surface (A12)	e (A11)		ark Surfa	ce (F6) rface (F7)		Red Parent Material (TF2)
	ucky Mineral (S1)		Redox D				Very Shallow Dark Surface (TF12) Other (Explain in Remarks)
	lucky Peat or Peat (S2) (LRR G, H)			ssions (F1	6)	³ Indicators of hydrophytic vegetation and
	cky Peat or Peat (S				3 of LRR	1997 - 1997 - 1997 - 1997 - 1997 - 1997 - 1997 - 1997 - 1997 - 1997 - 1997 - 1997 - 1997 - 1997 - 1997 - 1997 -	wetland hydrology must be present,
							unless disturbed or problematic.
	ayer (if present):						
Type:	/						. /
Depth (inc	hes):						Hydric Soil Present? Yes No
Remarks:							
HYDROLOG	GY			- All to water			
	rology Indicators:						
	ators (minimum of o		ock all that apply				Sacandan Indiantes (mining of the second
	Vater (A1)	no required, en	Salt Crust (Secondary Indicators (minimum of two required)
	er Table (A2)		Aquatic Inve		(P12)		Surface Soil Cracks (B6)
Saturation			Hydrogen S		- 12		Sparsely Vegetated Concave Surface (B8)
Water Ma			Dry-Season				Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3)
	Deposits (B2)		Oxidized Rt			Roote (
Drift Depo			(where no			9 110013 (Crayfish Burrows (C8)
Algal Mat	or Crust (B4)		Presence of		Iron (C4)		Saturation Visible on Aerial Imagery (C9)
Iron Depo	osits (B5)		Thin Muck S				Geomorphic Position (D2)
Inundation	n Visible on Aerial I	magery (B7)	Other (Expla		20		FAC-Neutral Test (D5)
	ained Leaves (B9)		~ · ·		0		Frost-Heave Hummocks (D7) (LRR F)
Field Observa	ations:					1	
Surface Water	Present? Ye	es No	Depth (incl	nes):			
Water Table P	Present? Yo		Depth (inch				1
Saturation Pre	sent? Ye		Depth (inch				and Hydrology Present? Yes No
(includes capil				3 570	1		We have a state of the second state of the sec
Describe Reco	orded Data (stream	gauge, monitori	ng well, aerial ph	iotos, pre	vious inspe	ections), i	if available:
Damada							
Remarks:							

Project/Site: Davis	City/County:	1 Paso	_ Sampling Date: <u>5/23/23</u>
Applicant/Owner: Davis		State: CO	Sampling Point: W6-WE
Investigator(s): Jon Dauzvardis	& GrantGUMESection, Towns	hip, Range: 533/34, T12	S,RGYW
Landform (hillslope, terrace, etc.): Stream		ncave, convex, none): _Con	cave Slope (%): 0-3
Subregion (LRR):	Lat: 38,961883	N Long: -104. 543	390° W Datum: WG5 84
Soil Map Unit Name: 19- Columbine Spe	welly syndy log m. 0-392 5		011500
Are climatic / hydrologic conditions on the site	typical for this time of year? Yes	No (If no, explain in	Remarks.)
Are Vegetation, Soil, or Hydro	logy significantly disturbed?	Are "Normal Circumstances"	
Are Vegetation, Soil, or Hydro	logy naturally problematic?	(If needed, explain any answ	State and a state of the state
SUMMARY OF FINDINGS - Attach	site map showing sampling p		
Hydrophytic Vegetation Present? Ye	No No		7
Hydric Soil Present? Ye	No Is the Sa	ampled Area	/*
Wetland Hydrology Present? Ye	s No within a	Wetland? Yes	No
Remarks:		A	
		1	
*			
VEGETATION – Use scientific nam	es of plants.		xi

Tree Stratum (Plot size:)	Absolute % Cover	Dominant Indicator Species? Status	Dominance Test worksheet:
1			Number of Dominant Species That Are OBL, FACW, or FAC
2			(excluding FAC-): (A)
3			Total Number of Dominant
4			Species Across All Strata: (B)
Sapling/Shrub Stratum (Plot size:)		= Total Cover	Percent of Dominant Species
1. Salix exigua	P	N OBL	That Are OBL, FACW, or FAC: (A/B)
2			Prevalence Index worksheet:
3			Total % Cover of:Multiply by:
4			OBL species x 1 =
5			FACW species x 2 =
		= Total Cover	FAC species x 3 =
Herb Stratum (Plot size:)	100	11	FACU species x 4 =
1. Juncus balticus	60	Y OBL	UPL species x 5 =
2. Eleocharis Palustris	20	Y OBL	Column Totals: (A) (B)
3. <u>Carex praegracilis</u> 4. Typha angostitolia	5	Y FACW	Prevalence Index = B/A =
-			Hydrophytic Vegetation Indicators:
6			1 - Rapid Test for Hydrophytic Vegetation
7			2 - Dominance Test is >50%
8			3 - Prevalence Index is ≤3.0 ¹
9			 4 - Morphological Adaptations¹ (Provide supporting data in Remarks or on a separate sheet)
10	AF		Problematic Hydrophytic Vegetation ¹ (Explain)
Woody Vine Stratum (Plot size:)		= Total Cover	¹ Indicators of hydric soil and wetland hydrology must
1			be present, unless disturbed or problematic.
2	100		Hydrophytic
% Bare Ground in Herb Stratum	100 =	Total Cover	Vegetation Present? Yes No
Remarks:			

US Army Corps of Engineers

Sampling Point: WG-WET

Depth Matrix	Redox Features	
(inches) Color (moist) %	Color (moist) % Type ¹ Loc ²	Texture Remarks
0-18 10YR5/1 100		Clayer Sand Saturated
		- tegan and and allo
Hydric Soil Indicators: (Applicable to al	I=Reduced Matrix, CS=Covered or Coated Sand	A CONTRACTOR OF
		Indicators for Problematic Hydric Soils ³ :
Histosol (A1)	Sandy Gleyed Matrix (S4)	1 cm Muck (A9) (LRR I, J)
Histic Epipedon (A2)	Sandy Redox (S5)	Coast Prairie Redox (A16) (LRR F, G, H)
Black Histic (A3)	Stripped Matrix (S6)	Dark Surface (S7) (LRR G)
Hydrogen Sulfide (A4) Stratified Lavers (A5) (LPR 5)	Loamy Mucky Mineral (F1)	High Plains Depressions (F16)
Stratified Layers (A5) (LRR F) 1 cm Muck (A9) (LRR F, G, H)	Loamy Gleyed Matrix (F2)	(LRR H outside of MLRA 72 & 73)
Depleted Below Dark Surface (A11)	 Depleted Matrix (F3) Redox Dark Surface (F6) 	Reduced Vertic (F18)
Thick Dark Surface (A12)	Depleted Dark Surface (F6)	Red Parent Material (TF2) Very Shallow Dark Surface (TF12)
Sandy Mucky Mineral (S1)	Redox Depressions (F8)	Other (Explain in Remarks)
2.5 cm Mucky Peat or Peat (S2) (LRR		³ Indicators of hydrophytic vegetation and
5 cm Mucky Peat or Peat (S3) (LRR F)		wetland hydrology must be present,
		unless disturbed or problematic.
Restrictive Layer (if present):		
Туре:		
Depth (inches):		Hydric Soil Present? Yes No
		Hydric Soil Present? Yes No
Depth (inches): Remarks:		Hydric Soil Present? Yes No
		Hydric Soil Present? Yes <u>V</u> No
		Hydric Soil Present? Yes No
Remarks:		Hydric Soil Present? Yes No
Remarks:		Hydric Soil Present? Yes No
Remarks: IYDROLOGY Wetland Hydrology Indicators:		Hydric Soil Present? Yes No
Remarks: IYDROLOGY Wetland Hydrology Indicators: Primary Indicators (minimum of one require	d; check all that apply)	Hydric Soil Present? Yes No
Remarks: YDROLOGY Wetland Hydrology Indicators: Primary Indicators (minimum of one require Surface Water (A1)	nd; check all that apply) Salt Crust (B11)	
Remarks: YDROLOGY Wetland Hydrology Indicators: Primary Indicators (minimum of one require Surface Water (A1) High Water Table (A2)		Secondary Indicators (minimum of two required)
Remarks: YDROLOGY Wetland Hydrology Indicators: Primary Indicators (minimum of one require Surface Water (A1)	Salt Crust (B11)	Secondary Indicators (minimum of two required) Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8)
Remarks: YDROLOGY Wetland Hydrology Indicators: Primary Indicators (minimum of one require Surface Water (A1) High Water Table (A2)	Salt Crust (B11) Aquatic Invertebrates (B13)	Secondary Indicators (minimum of two required) Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10)
Remarks: WDROLOGY Wetland Hydrology Indicators: Primary Indicators (minimum of one require Surface Water (A1) High Water Table (A2) Saturation (A3)	 Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) 	Secondary Indicators (minimum of two required) Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3)
Remarks: WDROLOGY Wetland Hydrology Indicators: Primary Indicators (minimum of one require Surface Water (A1) High Water Table (A2) Saturation (A3) Water Marks (B1)	 Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) 	 <u>Secondary Indicators (minimum of two required)</u> Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) s (C3) (where tilled)
Remarks: YDROLOGY Wetland Hydrology Indicators: Primary Indicators (minimum of one require Surface Water (A1) High Water Table (A2) Saturation (A3) Water Marks (B1) Sediment Deposits (B2)	 Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living Root (where not tilled) 	Secondary Indicators (minimum of two required) Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) s (C3) (where tilled) Crayfish Burrows (C8)
Remarks: Wetland Hydrology Indicators: Primary Indicators (minimum of one require Surface Water (A1) High Water Table (A2) Saturation (A3) Water Marks (B1) Sediment Deposits (B2) Drift Deposits (B3)	 Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living Root 	Secondary Indicators (minimum of two required) Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrows (C8) Saturation Visible on Aerial Imagery (C9)
Remarks: Wetland Hydrology Indicators: Primary Indicators (minimum of one require Surface Water (A1) High Water Table (A2) Saturation (A3) Water Marks (B1) Sediment Deposits (B2) Drift Deposits (B3) Algal Mat or Crust (B4) Iron Deposits (B5)	 Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living Root (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7) 	Secondary Indicators (minimum of two required) Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrows (C8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2)
Remarks: YDROLOGY Wetland Hydrology Indicators: Primary Indicators (minimum of one require Surface Water (A1) High Water Table (A2) Saturation (A3) Water Marks (B1) Sediment Deposits (B2) Drift Deposits (B3) Algal Mat or Crust (B4)	 Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living Root (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7) 	Secondary Indicators (minimum of two required) Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrows (C8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2) FAC-Neutral Test (D5)
Remarks: WDROLOGY Wetland Hydrology Indicators: Primary Indicators (minimum of one require Surface Water (A1) High Water Table (A2) Saturation (A3) Water Marks (B1) Sediment Deposits (B2) Drift Deposits (B3) Algal Mat or Crust (B4) Iron Deposits (B5) Inundation Visible on Aerial Imagery (B Water-Stained Leaves (B9)	 Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living Root (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7) 	Secondary Indicators (minimum of two required) Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrows (C8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2)
Remarks: YDROLOGY Wetland Hydrology Indicators: Primary Indicators (minimum of one require Surface Water (A1) High Water Table (A2) Saturation (A3) Water Marks (B1) Sediment Deposits (B2) Drift Deposits (B3) Algal Mat or Crust (B4) Iron Deposits (B5) Inundation Visible on Aerial Imagery (B Water-Stained Leaves (B9) Field Observations:	 Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living Root (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7) Other (Explain in Remarks) 	Secondary Indicators (minimum of two required) Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrows (C8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2) FAC-Neutral Test (D5)
Remarks: WDROLOGY Wetland Hydrology Indicators: Primary Indicators (minimum of one require Surface Water (A1) High Water Table (A2) Saturation (A3) Water Marks (B1) Sediment Deposits (B2) Drift Deposits (B3) Algal Mat or Crust (B4) Iron Deposits (B5) Inundation Visible on Aerial Imagery (B Water-Stained Leaves (B9) Field Observations: Surface Water Present? Yes	 Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living Root (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7) Other (Explain in Remarks) No Depth (inches):	Secondary Indicators (minimum of two required) Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrows (C8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2) FAC-Neutral Test (D5)
Remarks: YDROLOGY Wetland Hydrology Indicators: Primary Indicators (minimum of one require V Surface Water (A1) V High Water Table (A2) Saturation (A3) Water Marks (B1) Sediment Deposits (B2) Drift Deposits (B3) Algal Mat or Crust (B4) Iron Deposits (B5) Inundation Visible on Aerial Imagery (B Water-Stained Leaves (B9) Field Observations: Surface Water Present? Yes V Water Table Present? Yes V	 Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living Root (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7) Other (Explain in Remarks) 	Secondary Indicators (minimum of two required)Surface Soil Cracks (B6)Sparsely Vegetated Concave Surface (B8)Drainage Patterns (B10)Oxidized Rhizospheres on Living Roots (C3) s (C3) (where tilled)Crayfish Burrows (C8)Saturation Visible on Aerial Imagery (C9)Geomorphic Position (D2)FAC-Neutral Test (D5)Frost-Heave Hummocks (D7) (LRR F)
Remarks: WDROLOGY Wetland Hydrology Indicators: Primary Indicators (minimum of one require Surface Water (A1) High Water Table (A2) Saturation (A3) Water Marks (B1) Sediment Deposits (B2) Drift Deposits (B3) Algal Mat or Crust (B4) Iron Deposits (B5) Inundation Visible on Aerial Imagery (B Water-Stained Leaves (B9) Field Observations: Surface Water Present? Yes Water Table Present? Yes Saturation Present? Yes	 Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living Root (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7) Other (Explain in Remarks) 	Secondary Indicators (minimum of two required) Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrows (C8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2) FAC-Neutral Test (D5)
Remarks:	 Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living Root (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7) Other (Explain in Remarks) No Depth (inches): We We	Secondary Indicators (minimum of two required) Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Coxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrows (C8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2) FAC-Neutral Test (D5) Frost-Heave Hummocks (D7) (LRR F)
Remarks:	 Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living Root (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7) Other (Explain in Remarks) 	Secondary Indicators (minimum of two required) Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrows (C8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2) FAC-Neutral Test (D5) Frost-Heave Hummocks (D7) (LRR F)
Remarks:	 Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living Root (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7) Other (Explain in Remarks) No Depth (inches): We We	Secondary Indicators (minimum of two required) Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrows (C8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2) FAC-Neutral Test (D5) Frost-Heave Hummocks (D7) (LRR F)
Remarks:	 Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living Root (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7) Other (Explain in Remarks) No Depth (inches): We We	Secondary Indicators (minimum of two required) Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrows (C8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2) FAC-Neutral Test (D5) Frost-Heave Hummocks (D7) (LRR F)
Remarks:	 Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living Root (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7) Other (Explain in Remarks) No Depth (inches): We We	Secondary Indicators (minimum of two required) Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrows (C8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2) FAC-Neutral Test (D5) Frost-Heave Hummocks (D7) (LRR F)

Project/Site:	City/County: <u>El Paso</u> Sampling Date: <u>5/23/23</u>
Applicant/Owner:_ Davis	State: <u>CO</u> Sampling Point: <u>JN 6-UPL</u>
Investigator(s): Jan Dauzvardis + Grant Gurnée	_ Section, Township, Range: <u>53334, T125, R64W</u>
Landform (hillslope, terrace, etc.): <u>tervace</u>	Local relief (concave, convex, none): <u>Con Cave</u> Slope (%): 0-3%
Subregion (LRR): Lat: 3	8.961833°N Long: -104.543390°W Datum: WGS 84
Soil Map Unit Name: 19-Columbing growelly sandy le	aum, Q-390 slopes NWI classification:N/A
Are climatic / hydrologic conditions on the site typical for this time of y	year? Yes V No (If no, explain in Remarks.)
Are Vegetation, Soil, or Hydrology significant	ly disturbed? Are "Normal Circumstances" present? Yes No
Are Vegetation, Soil, or Hydrology naturally p	roblematic? (If needed, explain any answers in Remarks.)
SUMMARY OF FINDINGS – Attach site map showin	g sampling point locations, transects, important features, etc.
Hydrophytic Vegetation Present? Yes No	4
Hydric Soil Present? Yes No	- Is the Sampled Area
Wetland Hydrology Present? Yes No	within a Wetland? Yes No
Remarks:	
	/
М. .	
VEGETATION – Use scientific names of plants.	10

<u>Tree Stratum</u> (Plot size:) 1		Dominant Species?		Dominance Test workshow Number of Dominant Spec That Are OBL, FACW, or F	ties	
2				(excluding FAC-):	(2)/3	(A)
3				Total Number of Dominant		
4				Species Across All Strata:		(B)
Sapling/Shrub Stratum (Plot size:) 1)		= Total Cov	er	Percent of Dominant Spec That Are OBL, FACW, or F		_ (A/B)
		d a ann an a nn		Prevalence Index worksh	neet:	
2				Total % Cover of.	Multiply by:	
4				OBL species		
т б				FACW species	x 2 =	
		= Total Cov		FAC species		
Herb Stratum (Plot size:)	1.01	- Total Cov		FACU species	x 4 =	
1. Bromus inermis	10	Y	UPL	UPL species	x 5 =	
2. Pascopyrun smithii	70	Y	VPL	Column Totals:	2	
3				Prevalence Index [,] =	B/A =	
4				Hydrophytic Vegetation I		
5				1 - Rapid Test for Hyd		
6·				2 - Dominance Test is		
7				3 - Prevalence Index is		
8				4 - Morphological Ada		pointoga
9				data in Remarks or	on a separate shee	t)
10				Problematic Hydrophy	tic Vegetation ¹ (Exp	lain)
Woody Vine Stratum (Plot size:)		= Total Cov	er	¹ Indicators of hydric soil an be present, unless disturbe		/ must
1						
2	100	= Total Cov	er	Hydrophytic Vegetation Present? Yes _	No	
Remarks:	201201.508				¥	

Sampling Point: W____PL

Depth Matrix	Redox Features			icators.)
	Color (moist) % Type ¹	Loc ²	Texture	Remarks
0-18 ICVR-312 100		Sak	olyLoam	DN
			ergennin	
1-1-1-1-1-1-1-1-1-1-1-1-1-1-1-1-1-1-1-				
· · · · · · · · · · · · · · · · · · ·				
		· · · · · · · · · · · · · · · · · · ·		
¹ Type: C=Concentration, D=Depletion, RM=Red	uced Matrix. CS=Covered or Coat	ed Sand Grains	² Location:	PL=Pore Lining, M=Matrix.
Hydric Soil Indicators: (Applicable to all LRR	s, unless otherwise noted.)			oblematic Hydric Soils ³ :
Histosol (A1)	Sandy Gleyed Matrix (S4)			(9) (LRR I, J)
Histic Epipedon (A2)	Sandy Redox (S5)	-		Redox (A16) (LRR F, G, H)
Black Histic (A3)	Stripped Matrix (S6)	-		(S7) (LRR G)
Hydrogen Sulfide (A4)	Loamy Mucky Mineral (F1)			epressions (F16)
Stratified Layers (A5) (LRR F)	Loamy Gleyed Matrix (F2)			utside of MLRA 72 & 73)
1 cm Muck (A9) (LRR F, G, H)	Depleted Matrix (F3)		Reduced Ver	
Depleted Below Dark Surface (A11)	Redox Dark Surface (F6)		Red Parent M	laterial (TF2)
Thick Dark Surface (A12)	Depleted Dark Surface (F7) _	Very Shallow	Dark Surface (TF12)
Sandy Mucky Mineral (S1)	Redox Depressions (F8)	-		n in Remarks)
2.5 cm Mucky Peat or Peat (S2) (LRR G, H)		2200 T		ophytic vegetation and
5 cm Mucky Peat or Peat (S3) (LRR F)	(MLRA 72 & 73 of LRF	R H)	그는 것이 안 것 같은 것이 없어야 한다.	logy must be present,
Restrictive Layer (if present):			unless disturt	ed or problematic.
Turn				
Depth (inches):			1919-12012521	
	8	H	ydric Soil Prese	nt? Yes No
Remarks:				
				5 1 2
	2			
TYDROLOGY		Marriel Contraction	Ť.	
Wotland Hudrology Indicators				*
Wetland Hydrology Indicators:				•
Primary Indicators (minimum of one required; che	and the second state of th		Secondary Indi	cators (minimum of two required
Primary Indicators (minimum of one required; che Surface Water (A1)	Salt Crust (B11)		Surface So	il Cracks (B6)
Primary Indicators (minimum of one required; che Surface Water (A1) High Water Table (A2)	Salt Crust (B11) Aquatic Invertebrates (B13)		Surface So Sparsely V	il Cracks (B6) egetated Concave Surface (B8)
Primary Indicators (minimum of one required; che Surface Water (A1) High Water Table (A2) Saturation (A3)	Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1)		Surface So	il Cracks (B6) egetated Concave Surface (B8)
Primary Indicators (minimum of one required; che Surface Water (A1) High Water Table (A2) Saturation (A3) Water Marks (B1)	Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2)		Surface So Sparsely V Drainage P	il Cracks (B6) egetated Concave Surface (B8)
Primary Indicators (minimum of one required; che Surface Water (A1) High Water Table (A2) Saturation (A3) Water Marks (B1) Sediment Deposits (B2)	Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1)		Surface So Sparsely V Drainage P	il Cracks (B6) egetated Concave Surface (B8) atterns (B10) hizospheres on Living Roots (C3
Primary Indicators (minimum of one required; che Surface Water (A1) High Water Table (A2) Saturation (A3) Water Marks (B1) Sediment Deposits (B2) Drift Deposits (B3)	Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2)		Surface So Sparsely V Drainage P Oxidized R	il Cracks (B6) egetated Concave Surface (B8) atterns (B10) hizospheres on Living Roots (C3 lled)
Primary Indicators (minimum of one required; che	Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Liv	ing Roots (C3)	Surface So Sparsely V Drainage P Oxidized R (where ti Crayfish Bu	il Cracks (B6) egetated Concave Surface (B8) atterns (B10) hizospheres on Living Roots (C3 lled)
Primary Indicators (minimum of one required; che Surface Water (A1) High Water Table (A2) Saturation (A3) Water Marks (B1) Sediment Deposits (B2) Drift Deposits (B3) Algal Mat or Crust (B4) Iron Deposits (B5)	Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Liv (where not tilled)	ing Roots (C3)	Surface So Sparsely V Drainage P Oxidized R Oxidized R Crayfish Bu Saturation V	il Cracks (B6) egetated Concave Surface (B8) atterns (B10) hizospheres on Living Roots (C3 Iled) irrows (C8)
Primary Indicators (minimum of one required; che	Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Liv (where not tilled) Presence of Reduced Iron (C4)	ing Roots (C3)	Surface So Sparsely V Drainage P Oxidized R Oxidized R Crayfish Bu Saturation V	il Cracks (B6) egetated Concave Surface (B8) atterns (B10) hizospheres on Living Roots (C3 lled) irrøws (Č8) Visible on Aerial Imagery (C9) c Position (D2)
Primary Indicators (minimum of one required; che	Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Liv (where not tilled) Presence of Reduced Iron (C4 Thin Muck Surface (C7)	ing Roots (C3)	Surface So Sparsely V Drainage P Oxidized R Crayfish Bu Crayfish Bu Saturation V Geomorphi FAC-Neutra	il Cracks (B6) egetated Concave Surface (B8) atterns (B10) hizospheres on Living Roots (C3 lled) irrøws (Č8) Visible on Aerial Imagery (C9) c Position (D2)
Primary Indicators (minimum of one required; che	Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Liv (where not tilled) Presence of Reduced Iron (C4 Thin Muck Surface (C7)	ing Roots (C3)	Surface So Sparsely V Drainage P Oxidized R Crayfish Bu Crayfish Bu Saturation V Geomorphi FAC-Neutra	il Cracks (B6) egetated Concave Surface (B8) atterns (B10) hizospheres on Living Roots (C3 lled) urrows (Č8) Visible on Aerial Imagery (C9) c Position (D2) al Test (D5)
Primary Indicators (minimum of one required; che	Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Liv (where not tilled) Presence of Reduced Iron (C4 Thin Muck Surface (C7)	4)	Surface So Sparsely V Drainage P Oxidized R Crayfish Bu Crayfish Bu Saturation V Geomorphi FAC-Neutra	il Cracks (B6) egetated Concave Surface (B8) atterns (B10) hizospheres on Living Roots (C3 lled) urrows (Č8) Visible on Aerial Imagery (C9) c Position (D2) al Test (D5)
Primary Indicators (minimum of one required; che Surface Water (A1) High Water Table (A2) Saturation (A3) Water Marks (B1) Sediment Deposits (B2) Drift Deposits (B3) Algal Mat or Crust (B4) Iron Deposits (B5) Inundation Visible on Aerial Imagery (B7) Water-Stained Leaves (B9)	Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Liv (where not tilled) Presence of Reduced Iron (C4 Thin Muck Surface (C7) Other (Explain in Remarks)	(C3)	Surface So Sparsely V Drainage P Oxidized R Crayfish Bu Crayfish Bu Saturation V Geomorphi FAC-Neutra	il Cracks (B6) egetated Concave Surface (B8) atterns (B10) hizospheres on Living Roots (C3 lled) urrows (Č8) Visible on Aerial Imagery (C9) c Position (D2) al Test (D5)
Primary Indicators (minimum of one required; che Surface Water (A1) High Water Table (A2) Saturation (A3) Water Marks (B1) Sediment Deposits (B2) Drift Deposits (B3) Algal Mat or Crust (B4) Iron Deposits (B5) Inundation Visible on Aerial Imagery (B7) Water-Stained Leaves (B9)	Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Liv (where not tilled) Presence of Reduced Iron (C4 Thin Muck Surface (C7) Other (Explain in Remarks) Depth (inches):	4)	Surface So Sparsely V Drainage P Oxidized R (where ti Crayfish Bu Saturation V Geomorphi FAC-Neutra Frost-Heav	il Cracks (B6) egetated Concave Surface (B8) atterns (B10) hizospheres on Living Roots (C3 lied) urrøws (C8) Visible on Aerial Imagery (C9) c Position (D2) al Test (D5) e Hummocks (D7) (LRR F)
Primary Indicators (minimum of one required; che	Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Liv (where not tilled) Presence of Reduced Iron (C4 Thin Muck Surface (C7) Other (Explain in Remarks) Depth (inches): Depth (inches): Depth (inches):	4) Wetland F	Surface So Sparsely V Drainage P Oxidized R (where ti Crayfish Bu Saturation V Geomorphi FAC-Neutra Frost-Heav	il Cracks (B6) egetated Concave Surface (B8) atterns (B10) hizospheres on Living Roots (C3 lied) urrøws (C8) Visible on Aerial Imagery (C9) c Position (D2) al Test (D5) e Hummocks (D7) (LRR F)
Primary Indicators (minimum of one required; che Surface Water (A1) High Water Table (A2) Saturation (A3) Water Marks (B1) Sediment Deposits (B2) Drift Deposits (B3) Algal Mat or Crust (B4) Iron Deposits (B5) Inundation Visible on Aerial Imagery (B7) Water-Stained Leaves (B9) Field Observations: Surface Water Present? Yes No Water Table Present? Yes No Saturation Present? Yes No	Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Liv (where not tilled) Presence of Reduced Iron (C4 Thin Muck Surface (C7) Other (Explain in Remarks) Depth (inches): Depth (inches): Depth (inches):	4) Wetland F	Surface So Sparsely V Drainage P Oxidized R (where ti Crayfish Bu Saturation V Geomorphi FAC-Neutra Frost-Heav	il Cracks (B6) egetated Concave Surface (B8) atterns (B10) hizospheres on Living Roots (C3 lied) urrøws (C8) Visible on Aerial Imagery (C9) c Position (D2) al Test (D5) e Hummocks (D7) (LRR F)
Primary Indicators (minimum of one required; che	Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Liv (where not tilled) Presence of Reduced Iron (C4 Thin Muck Surface (C7) Other (Explain in Remarks) Depth (inches): Depth (inches): Depth (inches):	4) Wetland F	Surface So Sparsely V Drainage P Oxidized R (where ti Crayfish Bu Saturation V Geomorphi FAC-Neutra Frost-Heav	il Cracks (B6) egetated Concave Surface (B8) atterns (B10) hizospheres on Living Roots (C3 lied) urrøws (C8) Visible on Aerial Imagery (C9) c Position (D2) al Test (D5) e Hummocks (D7) (LRR F)
Primary Indicators (minimum of one required; che	Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Liv (where not tilled) Presence of Reduced Iron (C4 Thin Muck Surface (C7) Other (Explain in Remarks) Depth (inches): Depth (inches): Depth (inches):	4) Wetland F	Surface So Sparsely V Drainage P Oxidized R Crayfish Bu Crayfish Bu Saturation V Geomorphi FAC-Neutra Frost-Heav	il Cracks (B6) egetated Concave Surface (B8) atterns (B10) hizospheres on Living Roots (C3 lied) urrøws (C8) Visible on Aerial Imagery (C9) c Position (D2) al Test (D5) e Hummocks (D7) (LRR F)
Primary Indicators (minimum of one required; che	Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Liv (where not tilled) Presence of Reduced Iron (C4 Thin Muck Surface (C7) Other (Explain in Remarks) Depth (inches): Depth (inches): Depth (inches):	4) Wetland F	Surface So Sparsely V Drainage P Oxidized R Crayfish Bu Crayfish Bu Saturation V Geomorphi FAC-Neutra Frost-Heav	il Cracks (B6) egetated Concave Surface (B8) atterns (B10) hizospheres on Living Roots (C3 lied) urrøws (C8) Visible on Aerial Imagery (C9) c Position (D2) al Test (D5) e Hummocks (D7) (LRR F)
Primary Indicators (minimum of one required; che	Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Liv (where not tilled) Presence of Reduced Iron (C4 Thin Muck Surface (C7) Other (Explain in Remarks) Depth (inches): Depth (inches): Depth (inches):	4) Wetland F	Surface So Sparsely V Drainage P Oxidized R Crayfish Bu Crayfish Bu Saturation V Geomorphi FAC-Neutra Frost-Heav	il Cracks (B6) egetated Concave Surface (B8) atterns (B10) hizospheres on Living Roots (C3 lied) urrøws (C8) Visible on Aerial Imagery (C9) c Position (D2) al Test (D5) e Hummocks (D7) (LRR F)

Great Plains - Version 2.0

Project/Site: Davis	City/	County: <u>A Paso</u>	Sampling	g Date: 5/23/23
Applicant/Owner: Davis		St		Point: W7-WET
Investigator(s): Jon Dauzvardi	5 & GrantGumesecti	on, Township, Range [.]	T125, RG41	
Landform (hillslope, terrace, etc.): Stree		I relief (concave, convex, n	ione): <u>Concave</u>	Slope (%): 03
Subregion (LRR):	Lat: 38,96	21993° N Long: -	-104.543390° W	Datum: WGS 84
Soil Map Unit Name: 19-Columbin	e gravelly squale loan	0-392 slopes	NWI classification:	1/A
Are climatic / hydrologic conditions on the				
Are Vegetation, Soil, or Hy			ircumstances" present?	
Are Vegetation, Soil, or Hyperial of Hyperbolic Statements and the second statement of the	drology naturally problem		plain any answers in Rem	and the second se
SUMMARY OF FINDINGS - Atta	ich site map showing san	pling point location	s, transects, import	tant features, etc.
Hydrophytic Vegetation Present?	Yes No		100.000.000.000	and the second
Hydric Soil Present?	Yes No	Is the Sampled Area		
Wetland Hydrology Present?	Yes No	within a Wetland?	Yes No	
Remarks:			and the second second	

VEGETATION - Use scientific names of plants.

<u>Tree Stratum</u> (Plot size:) 1	Absolute % Cover	Dominant Indicator Species? Status	Dominance Test worksheet: Number of Dominant Species That Are OBL, FACW, or FAC
2			(excluding FAC-): (A)
3			Total Number of Dominant
4			Species Across All Strata: (B)
Sapling/Shrub Stratum (Plot size:) 1		= Total Cover	Percent of Dominant Species That Are OBL, FACW, or FAC: (A/B)
2			Prevalence Index worksheet:
3			Total % Cover of:Multiply by:
4			OBL species x 1 =
5			FACW species x 2 =
	A CONTRACTOR	= Total Cover	FAC species x 3 =
Herb Stratum (Plot size:)	- 10		FACU species x 4 =
1. Juncus balticus	60	Y OBL	UPL species x 5 =
2. Cavex Nebrascensis	40	Y OBL	Column Totals: (A) (B)
3			Prevalence Index = B/A =
5			Hydrophytic Vegetation Indicators:
6			1 - Rapid Test for Hydrophytic Vegetation
7			2 - Dominance Test is >50%
8			3 - Prevalence Index is ≤3.0 ¹
9			4 - Morphological Adaptations ¹ (Provide supporting data in Remarks or on a separate sheet)
10			Problematic Hydrophytic Vegetation ¹ (Explain)
Woody Vine Stratum (Plot size:) 1)		= Total Cover	¹ Indicators of hydric soil and wetland hydrology must be present, unless disturbed or problematic.
2		and the second second	
% Bare Ground in Herb Stratum	100/	= Total Cover	Hydrophytic Vegetation Present? Yes No
Remarks:			

r

Sampling Point: W7-WET

4	Redox Features	
	Color (moist)%Type ¹ o	c ² Texture Remarks
0-9 10YR3/1 100		SandyLeam
9-14 10YR3/1+3/2100		Sandylean Saturated @ 14"
14-18 104R2/1 100		Clay
		- Clay
Type: C=Concentration, D=Depletion, RM=Red Hydric Soil Indicators: (Applicable to all LRR:	uced Matrix, CS=Covered or Coated San	
Histosol (A1)		Indicators for Problematic Hydric Soils ³ :
Histosof (A1) Histic Epipedon (A2)	Sandy Gleyed Matrix (S4)	1 cm Muck (A9) (LRR I, J)
Black Histic (A3)	Sandy Redox (S5) Stripped Matrix (S6)	Coast Prairie Redox (A16) (LRR F, G, H)
Hydrogen Sulfide (A4)	Loamy Mucky Mineral (F1)	Dark Surface (S7) (LRR G)
Stratified Layers (A5) (LRR F)	Loamy Gleyed Matrix (F2)	High Plains Depressions (F16) (LRR H outside of MLRA 72 & 73)
1 cm Muck (A9) (LRR F, G, H)	Depleted Matrix (F3)	Reduced Vertic (F18)
Depleted Below Dark Surface (A11)	Redox Dark Surface (F6)	Red Parent Material (TF2)
Thick Dark Surface (A12)	Depleted Dark Surface (F7)	Very Shallow Dark Surface (TF12)
Sandy Mucky Mineral (S1)	Redox Depressions (F8)	Other (Explain in Remarks)
 2.5 cm Mucky Peat or Peat (S2) (LRR G, H) 5 cm Mucky Peat or Peat (S3) (LRR F) 		³ Indicators of hydrophytic vegetation and
	(MLRA 72 & 73 of LRR H)	wetland hydrology must be present,
estrictive Layer (if present):		unless disturbed or problematic.
Туре:		
Depth (inches):		
		Hydric Soil Present? Yes No
Remarks:	12	
Remarks:		
Remarks:	1. 1.	
emarks:	1) 1)	
	1)	
/DROLOGY	1) 1)	
PROLOGY /etland Hydrology Indicators:	ck all that apply)	Secondary Indicators (minimum of two required)
PROLOGY /etland Hydrology Indicators:		Secondary Indicators (minimum of two required)
/DROLOGY /etland Hydrology Indicators: rimary Indicators (minimum of one required; che	Salt Crust (B11)	Surface Soil Cracks (B6)
/DROLOGY /etland Hydrology Indicators: rimary Indicators (minimum of one required; che _ Surface Water (A1)	Salt Crust (B11) Aquatic Invertebrates (B13)	Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8)
/DROLOGY /etland Hydrology Indicators: rimary Indicators (minimum of one required; che Surface Water (A1) High Water Table (A2)	Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1)	 Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10)
Vetland Hydrology Indicators: rimary Indicators (minimum of one required; che _ Surface Water (A1) _ High Water Table (A2) _ Saturation (A3)	Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2)	 Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3)
/DROLOGY /etland Hydrology Indicators: rimary Indicators (minimum of one required; che Surface Water (A1) High Water Table (A2) Saturation (A3) Water Marks (B1)	Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living Rod	 Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled)
/DROLOGY /etland Hydrology Indicators: rimary Indicators (minimum of one required; che Surface Water (A1) High Water Table (A2) Saturation (A3) Water Marks (B1) Sediment Deposits (B2)	 Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living Rod (where not tilled) 	 Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrows (C8)
/DROLOGY /etland Hydrology Indicators: rimary Indicators (minimum of one required; che Surface Water (A1) High Water Table (A2) Saturation (A3) Water Marks (B1) Sediment Deposits (B2) Drift Deposits (B3)	 Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living Rod (where not tilled) Presence of Reduced Iron (C4) 	 Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrows (C8) Saturation Visible on Aerial Imagery (C9)
Vetland Hydrology Indicators: rimary Indicators (minimum of one required; che Surface Water (A1) High Water Table (A2) Saturation (A3) Water Marks (B1) Sediment Deposits (B2) Drift Deposits (B3) Algal Mat or Crust (B4)	Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living Rod (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7)	 Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrows (C8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2)
/DROLOGY /etland Hydrology Indicators: rimary Indicators (minimum of one required; che 	 Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living Rod (where not tilled) Presence of Reduced Iron (C4) 	 Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrows (C8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2) FAC-Neutral Test (D5)
/DROLOGY /etland Hydrology Indicators: rimary Indicators (minimum of one required; che 	Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living Rod (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7)	 Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrows (C8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2)
YDROLOGY Vetland Hydrology Indicators: rimary Indicators (minimum of one required; che Surface Water (A1) High Water Table (A2) Saturation (A3) Water Marks (B1) Sediment Deposits (B2) Drift Deposits (B3) Algal Mat or Crust (B4) Iron Deposits (B5) Inundation Visible on Aerial Imagery (B7) Water-Stained Leaves (B9) eld Observations:	Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living Rod (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7)	 Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrows (C8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2) FAC-Neutral Test (D5)
YDROLOGY Vetland Hydrology Indicators: trimary Indicators (minimum of one required; che Surface Water (A1) High Water Table (A2) Saturation (A3) Water Marks (B1) Sediment Deposits (B2) Drift Deposits (B3) Algal Mat or Crust (B4) Iron Deposits (B5) Inundation Visible on Aerial Imagery (B7) Water-Stained Leaves (B9) vetace Water Present? Yes	Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living Rod (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7) Other (Explain in Remarks) Depth (inches):	 Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrows (C8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2) FAC-Neutral Test (D5)
YDROLOGY Vetland Hydrology Indicators: trimary Indicators (minimum of one required; cheminimary Indicators (Mater Marks (Mater Table Present? Surface Water (A1) High Water Table (A2) Saturation (A3) Water Marks (B1) Sediment Deposits (B2) Drift Deposits (B3) Algal Mat or Crust (B4) Iron Deposits (B5) Inundation Visible on Aerial Imagery (B7) Water-Stained Leaves (B9) Vetater Present? Yes No Vater Table Present? Yes No	Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living Rod (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7) Other (Explain in Remarks)	 Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrows (C8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2) FAC-Neutral Test (D5) Frost-Heave Hummocks (D7) (LRR F)
YDROLOGY Vetland Hydrology Indicators: Primary Indicators (minimum of one required; chematical constructions) Surface Water (A1) High Water Table (A2) Saturation (A3) Water Marks (B1) Sediment Deposits (B2) Drift Deposits (B3) Algal Mat or Crust (B4) Iron Deposits (B5) Inundation Visible on Aerial Imagery (B7) Water-Stained Leaves (B9) Veter Table Present? Yes No Autor Present? Yes No No aturation Present? Yes Yes No	Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living Rod (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7) Other (Explain in Remarks) Depth (inches):	Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) ots (C3) (where tilled) Crayfish Burrows (C8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2) FAC-Neutral Test (D5) Frost-Heave Hummocks (D7) (LRR F) Mo
YDROLOGY Vetland Hydrology Indicators: rimary Indicators (minimum of one required; chematric chematrichematric chematric chematric chematric chematric chematric chemat	Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living Rod (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7) Other (Explain in Remarks) Depth (inches):	Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) ots (C3) (where tilled) Crayfish Burrows (C8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2) FAC-Neutral Test (D5) Frost-Heave Hummocks (D7) (LRR F) Mo
YDROLOGY Wetland Hydrology Indicators: Primary Indicators (minimum of one required; chematicators (minimum of one required; chematicators) Surface Water (A1) High Water Table (A2) Saturation (A3) Water Marks (B1) Sediment Deposits (B2) Drift Deposits (B3) Algal Mat or Crust (B4) Iron Deposits (B5) Inundation Visible on Aerial Imagery (B7) Water-Stained Leaves (B9) ield Observations: urface Water Present? Yes No aturation Present? Yes No noludes capillary fringe) escribe Recorded Data (stream gauge, monitoring)	Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living Rod (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7) Other (Explain in Remarks) Depth (inches):	Surface Soil Cracks (B6)Sparsely Vegetated Concave Surface (B8)Drainage Patterns (B10)Oxidized Rhizospheres on Living Roots (C3) ots (C3) (where tilled)Crayfish Burrows (C8)Saturation Visible on Aerial Imagery (C9)Geomorphic Position (D2)FAC-Neutral Test (D5)Frost-Heave Hummocks (D7) (LRR F)
YDROLOGY Vetland Hydrology Indicators: Primary Indicators (minimum of one required; chematical content of the second cont	Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living Rod (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7) Other (Explain in Remarks) Depth (inches):	Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) ots (C3) (where tilled) Crayfish Burrows (C8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2) FAC-Neutral Test (D5) Frost-Heave Hummocks (D7) (LRR F) Mo
YDROLOGY Vetland Hydrology Indicators: rimary Indicators (minimum of one required; chemistry Indicators (Mater Table (A2) Surface Water (A1) High Water Table (A2) Saturation (A3) Water Marks (B1) Sediment Deposits (B2) Drift Deposits (B3) Algal Mat or Crust (B4) Iron Deposits (B5) Inundation Visible on Aerial Imagery (B7) Water-Stained Leaves (B9) eld Observations: urface Water Present? Yes No daturation Present? Yes No No aturation Present? Yes No No cludes capillary fringe) No	Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living Rod (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7) Other (Explain in Remarks) Depth (inches):	Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrows (C8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2) FAC-Neutral Test (D5) Frost-Heave Hummocks (D7) (LRR F)
YDROLOGY Vetland Hydrology Indicators: rimary Indicators (minimum of one required; chemistry Indicators (Mater Table (A2) Surface Water (A1) High Water Table (A2) Saturation (A3) Water Marks (B1) Sediment Deposits (B2) Drift Deposits (B3) Algal Mat or Crust (B4) Iron Deposits (B5) Inundation Visible on Aerial Imagery (B7) Water-Stained Leaves (B9) eld Observations: urface Water Present? Yes No daturation Present? Yes No No aturation Present? Yes No No cludes capillary fringe) No	Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living Rod (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7) Other (Explain in Remarks) Depth (inches):	Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) ots (C3) (where tilled) Crayfish Burrows (C8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2) FAC-Neutral Test (D5) Frost-Heave Hummocks (D7) (LRR F) Mo

Project/Site: Davis	City/County: <u>El Paso</u> Sampling Date: <u>5/23/23</u>
Applicant/Owner: Davis	State: <u>CO</u> Sampling Point: <u>W7-UPL</u>
Investigator(s): Jan Davzvardis + Grant Gurnée	Section, Township, Range: 533734, TI25, R64W
Landform (hillslope, terrace, etc.): <u>tervace</u>	
Subregion (LRR): 4	3.961883. N Long: -104. 543390° W Datum: WGS 84
Soil Map Unit Name: 19- Columbine gravely bandy bam, C	2-3% slupes NWI classification: N/A
Are climatic / hydrologic conditions on the site typical for this time of ye	ear? Yes No (If no, explain in Remarks.)
Are Vegetation, Soil, or Hydrology significantly	/ disturbed? Are "Normal Circumstances" present? Yes Ves No
Are Vegetation, Soil, or Hydrology naturally pro	oblematic? (If needed, explain any answers in Remarks.)
SUMMARY OF FINDINGS - Attach site map showing	g sampling point locations, transects, important features, etc.

Hydrophytic Vegetation Present? Hydric Soil Present? Wetland Hydrology Present?	Yes Yes Yes	Is the Sampled Area within a Wetland?	Yes	No	
Remarks:		 			
•					

VEGETATION – Use scientific names of plants.

Tree Stratum (Plot size:) 1		= Total Cover	Dominance Test worksheet: Number of Dominant Species That Are OBL, FACW, or FAC (excluding FAC-): Total Number of Dominant Species Across All Strata: Percent of Dominant Species That Are OBL, FACW, or FAC: (A)
1			Prevalence Index worksheet:
2			Total % Cover of: Multiply by:
3	8 <u>-1</u>		$\frac{1}{\text{OBL species}} = \frac{5}{x^{1}} = \frac{5}{x^{1}}$
4			FACW species x 2 =
5			FAC species x 2 =
Herb Stratum (Plot size:)		= Total Cover	FACU species $\underline{40}$ $x4 = \underline{160}$
1. <u>Civilum arvense</u>	40	V FACU	UPL species $10 \times 5 = 50$
2. Machaeranthera Fanacetitoli	45	VUPL	Column Totals: <u>5,2</u> (A) <u>220</u> (B)
3. America fruiticosa.	5	VOBL	110
4. Gevanium viscosissimum	5	VNT	Prevalence Index' = B/A =
5. Verbascum Thasars	6	U UPL	Hydrophytic Vegetation Indicators:
			1 - Rapid Test for Hydrophytic Vegetation
			2 - Dominance Test is >50%
7			3 - Prevalence Index is ≤3.0 ¹
8			4 - Morphological Adaptations ¹ (Provide supporting
9			data in Remarks or on a separate sheet)
10	60	= Total Cover	Problematic Hydrophytic Vegetation ¹ (Explain)
Woody Vine Stratum (Plot size:)			¹ Indicators of hydric soil and wetland hydrology must
1			be present, unless disturbed or problematic.
2			Hydrophytic
% Bare Ground in Herb Stratum	100	= Total Cover	Vegetation Present? Yes No
Remarks:			

US Army Corps of Engineers

Great Plains - Version 2.0

Sampling Point: W_7-UPL

Profile Description: (Desc	ribe to the depth nee	ded to document the indicator or	confirm the absence of indicators.)
Depth Mat		Redox Features	
(inches) Color (mois	t) % Co	lor (moist) % Type ¹	Loc ² Texture Remarks
0-6 [04R3	3/2		Sandy Clay Loam
6-12 10 YR3	3/2		Loamy Clay
17-10, TOVE 31	7.		Clav
1210 10/10/1			<u>Ctay</u>
Constanting of the Constant of the	ne state		
		ced Matrix, CS=Covered or Coated	
Hydric Soil Indicators: (Ap	oplicable to all LRRs,	unless otherwise noted.)	Indicators for Problematic Hydric Soils ³ :
Histosol (A1)		Sandy Gleyed Matrix (S4)	1 cm Muck (A9) (LRR I, J)
Histic Epipedon (A2)		Sandy Redox (S5)	Coast Prairie Redox (A16) (LRR F, G, H)
Black Histic (A3)		Stripped Matrix (S6)	Dark Surface (S7) (LRR G)
Hydrogen Sulfide (A4)		Loamy Mucky Mineral (F1)	High Plains Depressions (F16)
Stratified Layers (A5) (L		Loamy Gleyed Matrix (F2)	(LRR H outside of MLRA 72 & 73)
1 cm Muck (A9) (LRR F,		Depleted Matrix (F3)	Reduced Vertic (F18)
Depleted Below Dark Su Thick Dark Surface (A12		Redox Dark Surface (F6)	Red Parent Material (TF2)
Sandy Mucky Mineral (S		Depleted Dark Surface (F7) Redox Depressions (F8)	Very Shallow Dark Surface (TF12)
2.5 cm Mucky Peat or P		High Plains Depressions (F16)	Other (Explain in Remarks) ³ Indicators of hydrophytic vegetation and
5 cm Mucky Peat or Pea		(MLRA 72 & 73 of LRR H	
		(unless disturbed or problematic.
Restrictive Layer (if presen	nt):	NAMES OF THE INCOME.	
Type:			
	and the second second		Hydric Soil Present? Yes No
Remarks:	1		
Tremarka.			
HYDROLOGY			
Wetland Hydrology Indicat	ore	······	
		k all that each à	
Primary Indicators (minimum	i of one required; chec		Secondary Indicators (minimum of two required)
Surface Water (A1)		Salt Crust (B11)	Surface Soil Cracks (B6)
High Water Table (A2)		_ Aquatic Invertebrates (B13)	Sparsely Vegetated Concave Surface (B8)
Saturation (A3)		_ Hydrogen Sulfide Odor (C1)	Drainage Patterns (B10)
Water Marks (B1)	-	_ Dry-Season Water Table (C2)	Oxidized Rhizospheres on Living Roots (C3)
Sediment Deposits (B2)		Oxidized Rhizospheres on Living	
Drift Deposits (B3)		(where not tilled)	Crayfish Burrøws (C8)
Algal Mat or Crust (B4)	-	Presence of Reduced Iron (C4)	Saturation Visible on Aerial Imagery (C9)
Iron Deposits (B5)		_ Thin Muck Surface (C7)	Geomorphic Position (D2)
Inundation Visible on Ae	rial Imagery (B7)	Other (Explain in Remarks)	FAC-Neutral Test (D5)
Water-Stained Leaves (E	39)		Frost-Heave Hummocks (D7) (LRR F)
Field Observations:			
Surface Water Present?	Yes No	Depth (inches):	
Water Table Present?		Depth (inches):	
Saturation Present?	and the second sec	Depth (inches):	Wetland Hydrology Present? Yes No
(includes capillary fringe)			
Describe Recorded Data (str	eam gauge, monitorin	g well, aerial photos, previous inspec	ctions), if available:
Remarks:			

Project/Site: Davis	City/County: El Pas	O Sampling	Date: 5/23/23
Applicant/Owner: Davis		State: CO Sampling	Point: WO-WET
Investigator(s): Jon Dauzvardis & Grant	GUTHE Section, Township, Ran	ge: 533-3-1, T125, RG4U	/
Landform (hillslope, terrace, etc.): Stream Corridor	Local relief /concave, c	onvex, none): <u>Concave</u>	Slope (%):
Subregion (LRR):			Datum: WGS 84
Soil Map Unit Name: 29-Fluxagregtic Mapley	valls, nearly level	NWI classification:	EM1A
Are climatic / hydrologic conditions on the site typical for this	time of year? Yes No	(If no, explain in Remarks.)	
Are Vegetation, Soil, or Hydrology sig	inificantly disturbed? Are "	Normal Circumstances" present? Y	res No
Are Vegetation, Soil, or Hydrology na	turally problematic? (If nee	eded, explain any answers in Rema	rks.)
SUMMARY OF FINDINGS – Attach site map s	howing sampling point lo	cations, transects, import	ant features, etc.
		7	

Hydrophytic Vegetation Present? Hydric Soil Present? Wetland Hydrology Present?	Yes No Yes No Yes No	Is the Sampled Area within a Wetland?	Yes No
Remarks:		/	
		1	

VEGETATION - Use scientific names of plants.

Tree Stratum (Plot size:) 1)		Dominant Indicator Species? Status	Dominance Test worksheet: Number of Dominant Species That Are OBL, FACW, or FAC
2			(excluding FAC-):
3			Total Number of Dominant
4			Species Across All Strata: (B)
Sapling/Shrub Stratum (Plot size:)		= Total Cover	Percent of Dominant Species That Are OBL, FACW, or FAC: (A/B)
1			Prevalence Index worksheet:
2			Total % Cover of:Multiply by:
3		······································	OBL species x 1 =
4 5			FACW species x 2 =
		= Total Cover	FAC species x 3 =
Herb Stratum (Plot size:)			FACU species x 4 =
1. Couvex hebrascensis	50	Y OBL	UPL species x 5 =
2. Juncus balticus	25	Y OBL	Column Totals: (A) (B)
3. Typha angustitelia	10	Y OBL	
4. Grsjund grvenst	5	Y FACY	Prevalence Index = B/A =
5. Mentha aquaticy	10	Y OBL	Hydrophytic Vegetation Indicators:
6l			1 - Rapid Test for Hydrophytic Vegetation
7			2 - Dominance Test is >50%
8			3 - Prevalence Index is ≤3.0 ¹
9			4 - Morphological Adaptations ¹ (Provide supporting data in Remarks or on a separate sheet)
10			Problematic Hydrophytic Vegetation ¹ (Explain)
Woody Vine Stratum (Plot size:)	100	= Total Cover	¹ Indicators of hydric soil and wetland hydrology must
1			be present, unless disturbed or problematic.
2			Hydrophytic
% Bare Ground in Herb Stratum	100	= Total Cover	Vegetation Present? Yes No
Remarks:			

Sampling Point: W8-WET

Profile Description: (Descri	be to the depth ne	eded to docum	ent the i	ndicator o	or confirm	n the absence of indicators.)
Depth Matri			Features			
(inches) Color (moist)	<u>%</u> C	olor (moist)		Type ¹	_Loc ²	Texture Remarks
0-2 014				-		organic Matter
2-12 10YR2	1 100					SilfySand
12-18 107R2/1+4	1/ iau					Clay
have been been been been been been been be						and the providence of the second s
¹ Type: C=Concentration, D=I	Depletion, RM=Red	uced Matrix, CS	=Covered	or Coate	d Sand G	rains. ² Location: PL=Pore Lining, M=Matrix.
Hydric Soil Indicators: (App	plicable to all LRR	s, unless other	wise note	ed.)		Indicators for Problematic Hydric Soils ³ :
Histosol (A1)		Sandy G	leyed Ma	trix (S4)		1 cm Muck (A9) (LRR I, J)
Histic Epipedon (A2)		the second s	edox (S5			Coast Prairie Redox (A16) (LRR F, G, H)
Black Histic (A3)			Matrix (S			Dark Surface (S7) (LRR G)
V Hydrogen Sulfide (A4)			lucky Min			High Plains Depressions (F16)
Stratified Layers (A5) (LR			Bleyed Ma			(LRR H outside of MLRA 72 & 73)
1 cm Muck (A9) (LRR F, Depleted Below Dark Sur		the second se	I Matrix (F ark Surfa			Reduced Vertic (F18) Red Parent Material (TF2)
Thick Dark Surface (A12)		1.7757 C		rface (F7)		Very Shallow Dark Surface (TF12)
Sandy Mucky Mineral (S1			epression			Other (Explain in Remarks)
2.5 cm Mucky Peat or Pe				ssions (F	16)	³ Indicators of hydrophytic vegetation and
5 cm Mucky Peat or Peat	(S3) (LRR F)	(MLF	RA 72 & 7	3 of LRR	H)	wetland hydrology must be present,
						unless disturbed or problematic.
Restrictive Layer (if present	:):					
Depth (inches):		- E				Hydric Soil Present? Yes V No
Remarks:						
					170	
HYDROLOGY						
Wetland Hydrology Indicato	rs:					6.je.
Primary Indicators (minimum	of one required; che	eck all that apply)			Secondary Indicators (minimum of two required)
Surface Water (A1)		Salt Crust (B11)			Surface Soil Cracks (B6)
High Water Table (A2)		Aquatic Inv	ertebrate	s (B13)		Sparsely Vegetated Concave Surface (B8)
Saturation (A3)		Hydrogen S	Sulfide Oc	lor (C1)		Drainage Patterns (B10)
Vater Marks (B1)		Dry-Seasor	Water T	able (C2)		Oxidized Rhizospheres on Living Roots (C3)
Sediment Deposits (B2)		Oxidized R	hizospher	es on Livi	ng Roots	(C3) (where tilled)
V Drift Deposits (B3)		(where n	ot tilled)			Crayfish Burrows (C8)
Algal Mat or Crust (B4)		Presence o	f Reduce	d Iron (C4)	Saturation Visible on Aerial Imagery (C9)
Iron Deposits (B5)		Thin Muck	Surface (C7)		Geomorphic Position (D2)
Inundation Visible on Aer	ial Imagery (B7)	Other (Expl	ain in Re	marks)		FAC-Neutral Test (D5)
Water-Stained Leaves (B	9)					Frost-Heave Hummocks (D7) (LRR F)
Field Observations:	11		^	1	-	
Surface Water Present?	Yes No	Depth (inc	hes): 0	1	-	
Water Table Present?	Yes V No	Depth (inc	hes): <u>()</u>	(Surtag	2	. /
Saturation Present? (includes capillary fringe)	Yes No	Depth (inc	hes): <u>((</u>	surtag	Weth	and Hydrology Present? Yes No
Describe Recorded Data (stre	am gauge, monitori	ng well, aerial p	hotos, pre	evious insp	pections),	if available:
Remarks:						

Project/Site: Davis	City/County: <u>El Paso</u>	Sampling D	ate: <u>5/23/23</u>
Applicant/Owner: Davis	State	: <u>CO</u> _ Sampling Po	oint: W8-UPL
Investigator(s): Jan Dauzvardis + Grant Gurnée	Section, Township, Range: 533/3	4, T125, R64U	<u> </u>
Landform (hillslope, terrace, etc.): <u>tervace</u>	Local relief (concave, convex, non	e): concare	Slope (%): <u>Q-2. %</u>
Subregion (LRR): Lat:3	3.945566°N Long: -10	04.529015°W	Datum: WGS 84
Subregion (LRR): <u>G</u> Soil Map Unit Name: <u>29- Flu vaquentic H</u> uploquallis, n	early level.	NWI classification:	V/A
Are climatic / hydrologic conditions on the site typical for this time of ye		, explain in Remarks.)	
Are Vegetation, Soil, or Hydrology significantly	disturbed? Are "Normal Circ	umstances" present? Yes	s No
Are Vegetation, Soil, or Hydrology naturally pr	oblematic? (If needed, expla	in any answers in Remarks	s.)
SUMMARY OF FINDINGS – Attach site map showing	sampling point locations,	transects, importar	nt features, etc.

Hydrophytic Vegetation Present? Hydric Soil Present? Wetland Hydrology Present?	Yes Yes Yes	No No	Is the Sampled Area within a Wetland?	Yes	No	
Remarks:						
			· · · · · · · · · · · · · · · · · · ·			

VEGETATION - Use scientific names of plants.

Tree Stratum (Plot size:) 1 2			Dominance Test worksheet: Number of Dominant Species That Are OBL, FACW, or FAC (excluding FAC-):	(A)
3	-		Total Number of Dominant Species Across All Strata:	B)
4		= Total Cover	Percent of Dominant Species	A/B)
12			Prevalence Index worksheet:	-
23			Total % Cover of: Multiply by:	
4			OBL species x 1 =	
5	-		FACW species x 2 =	
		= Total Cover	FAC species x 3 =	
Herb Stratum (Plot size:)		- Total Cover	FACU species x 4 =	
1. Bosteloug avacilis	10	UPL	UPL species x 5 =	
2. Machaevanthara tanacetitalia	20	UPL	Column Totals: (A)	(B)
3. Artemisia trigida	10	VPL	Prevalence Index [,] = B/A =	
4			Hydrophytic Vegetation Indicators:	_
5			1 - Rapid Test for Hydrophytic Vegetation	
67			2 - Dominance Test is >50%	
7			3 - Prevalence Index is ≤3.0 ¹	
89			4 - Morphological Adaptations ¹ (Provide suppo data in Remarks or on a separate sheet)	rting
10			Problematic Hydrophytic Vegetation ¹ (Explain)	
Woody Vine Stratum (Plot size:) 1	40	= Total Cover	¹ Indicators of hydric soil and wetland hydrology mu be present, unless disturbed or problematic.	
2.			Hydrophytic	
% Bare Ground in Herb Stratum60	100	= Total Cover	Vegetation Present? Yes No	
Remarks:				

C	2		8	
Э	v	s	L	é

Sampling Point: W8-UPL

	Charles a second s	onfirm the absence of indicators.)
Depth <u>Matrix</u> (inches) Color (moist)	% Color (moist) % Type ¹ Lo	oc ² Texture Remarks
0-10 10YR3/2	100	
	100	Sandyloam
10-18 104R312 1		Sandy Gravelly Leam
	ion, RM=Reduced Matrix, CS=Covered or Coated Sa	and Grains. ² Location: PL=Pore Lining, M=Matrix.
Hydric Soil Indicators: (Applicabl	le to all LRRs, unless otherwise noted.)	Indicators for Problematic Hydric Soils ³ :
Histosol (A1)	Sandy Gleyed Matrix (S4)	1 cm Muck (A9) (LRR I, J)
Histic Epipedon (A2)	Sandy Redox (S5)	Coast Prairie Redox (A16) (LRR F, G, H)
Black Histic (A3)	Stripped Matrix (S6)	Dark Surface (S7) (LRR G)
Hydrogen Sulfide (A4) Stratified Lavers (A5) (LBB E)	Loamy Mucky Mineral (F1)	High Plains Depressions (F16)
Stratified Layers (A5) (LRR F) 1 cm Muck (A9) (LRR F, G, H)	Loamy Gleyed Matrix (F2) Depleted Matrix (F3)	(LRR H outside of MLRA 72 & 73) Reduced Vertic (F18)
Depleted Below Dark Surface (A		Red Parent Material (TF2)
Thick Dark Surface (A12)	Depleted Dark Surface (F7)	Very Shallow Dark Surface (TF12)
Sandy Mucky Mineral (S1)	Redox Depressions (F8)	Other (Explain in Remarks)
2.5 cm Mucky Peat or Peat (S2)		³ Indicators of hydrophytic vegetation and
5 cm Mucky Peat or Peat (S3) ((LRR F) (MLRA 72 & 73 of LRR H)	wetland hydrology must be present,
Profiling Low (16		unless disturbed or problematic.
Restrictive Layer (if present):		
Туре:		
Depth (inches):		Hydric Soil Present? Yes No
Remarks:		
HYDROLOGY		· •
Wetland Hydrology Indicators:		
Primary Indicators (minimum of one	28 31 8 76 3278 827	
	required; check all that apply)	Secondary Indicators (minimum of two required)
Surface Water (A1)	salt Crust (B11)	
Surface Water (A1) High Water Table (A2)		Secondary Indicators (minimum of two required) Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8)
	Salt Crust (B11) Aquatic Invertebrates (B13)	Surface Soil Cracks (B6)
High Water Table (A2)	Salt Crust (B11)	 Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8)
High Water Table (A2) Saturation (A3)	Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1)	 Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3)
High Water Table (A2) Saturation (A3) Water Marks (B1)	 Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) 	 Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3)
 High Water Table (A2) Saturation (A3) Water Marks (B1) Sediment Deposits (B2) 	 Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living Filteres (C2) 	 Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled)
 High Water Table (A2) Saturation (A3) Water Marks (B1) Sediment Deposits (B2) Drift Deposits (B3) Algal Mat or Crust (B4) Iron Deposits (B5) 	 Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living F (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7) 	 Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrøws (C8)
 High Water Table (A2) Saturation (A3) Water Marks (B1) Sediment Deposits (B2) Drift Deposits (B3) Algal Mat or Crust (B4) 	 Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living F (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7) 	 Surface Soil-Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrøws (C8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2) FAC-Neutral Test (D5)
 High Water Table (A2) Saturation (A3) Water Marks (B1) Sediment Deposits (B2) Drift Deposits (B3) Algal Mat or Crust (B4) Iron Deposits (B5) Inundation Visible on Aerial Ima Water-Stained Leaves (B9) 	 Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living F (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7) 	 Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrøws (C8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2)
 High Water Table (A2) Saturation (A3) Water Marks (B1) Sediment Deposits (B2) Drift Deposits (B3) Algal Mat or Crust (B4) Iron Deposits (B5) Inundation Visible on Aerial Ima Water-Stained Leaves (B9) Field Observations:	Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living F (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7) agery (B7) Other (Explain in Remarks)	 Surface Soil-Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrøws (C8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2) FAC-Neutral Test (D5)
 High Water Table (A2) Saturation (A3) Water Marks (B1) Sediment Deposits (B2) Drift Deposits (B3) Algal Mat or Crust (B4) Iron Deposits (B5) Inundation Visible on Aerial Ima Water-Stained Leaves (B9) Field Observations: Surface Water Present? Yes 	Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living F (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7) Other (Explain in Remarks)	 Surface Soil-Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrøws (C8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2) FAC-Neutral Test (D5)
 High Water Table (A2) Saturation (A3) Water Marks (B1) Sediment Deposits (B2) Drift Deposits (B3) Algal Mat or Crust (B4) Iron Deposits (B5) Inundation Visible on Aerial Ima Water-Stained Leaves (B9) Field Observations: Surface Water Present? Yes Water Table Present? Yes	 Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living F (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7) Other (Explain in Remarks) 	 Surface Soil-Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrøws (C8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2) FAC-Neutral Test (D5) Frost-Heave Hummocks (D7) (LRR F)
High Water Table (A2) Saturation (A3) Water Marks (B1) Sediment Deposits (B2) Drift Deposits (B3) Algal Mat or Crust (B4) Iron Deposits (B5) Inundation Visible on Aerial Ima Water-Stained Leaves (B9) Field Observations: Surface Water Present? Yes Water Table Present? Yes Saturation Present? Yes	Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living F (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7) Other (Explain in Remarks)	 Surface Soil-Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrøws (C8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2) FAC-Neutral Test (D5)
High Water Table (A2) Saturation (A3) Water Marks (B1) Sediment Deposits (B2) Drift Deposits (B3) Algal Mat or Crust (B4) Iron Deposits (B5) Inundation Visible on Aerial Ima Water-Stained Leaves (B9) Field Observations: Surface Water Present? Yes Water Table Present? Yes Saturation Present? Yes (includes capillary fringe)	 Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living F (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7) Other (Explain in Remarks) 	Surface Soil-Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrøws (C8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2) FAC-Neutral Test (D5) Frost-Heave Hummocks (D7) (LRR F) Wetland Hydrology Present? Yes No
 High Water Table (A2) Saturation (A3) Water Marks (B1) Sediment Deposits (B2) Drift Deposits (B3) Algal Mat or Crust (B4) Iron Deposits (B5) Inundation Visible on Aerial Ima Water-Stained Leaves (B9) Field Observations: Surface Water Present? Yes Water Table Present? Yes Saturation Present? Yes (includes capillary fringe) 	 Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living F (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7) Other (Explain in Remarks) 	Surface Soil-Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrøws (C8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2) FAC-Neutral Test (D5) Frost-Heave Hummocks (D7) (LRR F) Wetland Hydrology Present? Yes No
 High Water Table (A2) Saturation (A3) Water Marks (B1) Sediment Deposits (B2) Drift Deposits (B3) Algal Mat or Crust (B4) Iron Deposits (B5) Inundation Visible on Aerial Ima Water-Stained Leaves (B9) Field Observations: Surface Water Present? Yes Water Table Present? Yes (includes capillary fringe) Describe Recorded Data (stream gate) 	 Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living F (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7) Other (Explain in Remarks) 	Surface Soil-Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrøws (C8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2) FAC-Neutral Test (D5) Frost-Heave Hummocks (D7) (LRR F) Wetland Hydrology Present? Yes No
High Water Table (A2) Saturation (A3) Water Marks (B1) Sediment Deposits (B2) Drift Deposits (B3) Algal Mat or Crust (B4) Iron Deposits (B5) Inundation Visible on Aerial Ima Water-Stained Leaves (B9) Field Observations: Surface Water Present? Yes Water Table Present? Yes Saturation Present? Yes (includes capillary fringe)	 Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living F (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7) Other (Explain in Remarks) 	Surface Soil-Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrøws (C8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2) FAC-Neutral Test (D5) Frost-Heave Hummocks (D7) (LRR F) Wetland Hydrology Present? Yes No
 High Water Table (A2) Saturation (A3) Water Marks (B1) Sediment Deposits (B2) Drift Deposits (B3) Algal Mat or Crust (B4) Iron Deposits (B5) Inundation Visible on Aerial Ima Water-Stained Leaves (B9) Field Observations: Surface Water Present? Yes Water Table Present? Yes (includes capillary fringe) Describe Recorded Data (stream gate) 	 Salt Crust (B11) Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living F (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7) Other (Explain in Remarks) 	Surface Soil-Cracks (B6) Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrøws (C8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2) FAC-Neutral Test (D5) Frost-Heave Hummocks (D7) (LRR F) Wetland Hydrology Present? Yes No

Project/Site:Qavis	City/County: El Paso	Sampling Date: 5/23/23
Applicant/Owner: Davis	State:	00 110 217
Investigator(s): Jon Dauzvardis & Grant	10106 Section, Township, Range: 533-34	
Landform (hillslope, terrace, etc.): <u>Stream Corridor</u>	Local relief (concave, convex, none	~ ~ ~
Subregion (LRR):	Lat: <u>38 - 961983° N</u> Long: <u>-10</u>	4.543390° W Datum: WGS 84
Soil Map Unit Name: 29 - Fluxaquentic Hap	laguolls, nearly level ,	WI classification: N/A
Are climatic / hydrologic conditions on the site typical for this ti	me of year? Yes No (If no,	explain in Remarks.)
Are Vegetation, Soil, or Hydrology sign	ificantly disturbed? Are "Normal Circu	mstances" present? Yes V No
Are Vegetation, Soil, or Hydrology nate	urally problematic? (If needed, explain	any answers in Remarks.)
SUMMARY OF FINDINGS - Attach site map sh	owing sampling point locations.	ransects, important features, etc.
Hydrophytic Vegetation Present? Yes No Hydric Soil Present? Yes No Wetland Hydrology Present? Yes No	Is the Sampled Area within a Wetland?	Yes No
Remarks:	2	
	~	
· ,		

VEGETATION - Use scientific names of plants.

Tree Stratum (Plot size:) 1			
4			Species Across All Strata:
Sapling/Shrub Stratum (Plot size:) 1		= Total Cover	Percent of Dominant Species
2		· · · · · · · · · · · · · · · · · · ·	Prevalence Index worksheet:
3			Total % Cover of: Multiply by:
4			OBL species x 1 =
5.			FACW species x 2 =
		= Total Cover	FAC species x 3 =
Herb Stratum (Plot size:)	10	11 001	FACU species x 4 =
1. Juneus batticus	-te	Y OBL	UPL species x 5 =
2. Cavex praegracilis	5	Y HEV	Column Totals: (A) (B)
3. Distichlis Spicate	25	_Y_FAC	Prevalence Index = B/A =
5			Hydrophytic Vegetation Indicators:
6			1 - Rapid Test for Hydrophytic Vegetation
7			2 - Dominance Test is >50%
8			3 - Prevalence Index is ≤3.0 ¹
9			 4 - Morphological Adaptations¹ (Provide supporting data in Remarks or on a separate sheet)
10	-		 Problematic Hydrophytic Vegetation¹ (Explain)
	40	= Total Cover	¹ Indicators of hydric soil and wetland hydrology must
Woody Vine Stratum (Plot size:)			be present, unless disturbed or problematic.
1			- Ibudaaabadla /
Bare Ground in Herb Stratum	100	= Total Cover	_ Hydrophytic Vegetation Present? Yes No
Remarks:	8-1-1-1-1-1-1		

US Army Corps of Engineers

Sampling Point: M9-WET

Profile Descr	ription: (Describe	to the depth n	eeded to document the indic	ator or confirm	the absence of indicators.)
Depth	Matrix		Redox Features		
(inches)	Color (moist)	%(Color (moist) % Ty	pe ¹ Loc ²	Texture Remarks
0-10	104K311			>	nttylcam Saturated @ 12"
					1
	•••••••••••••••••				
and the second se	and the local division of the second division		duced Matrix, CS=Covered or (Coated Sand Gra	
Hydric Soil II	ndicators: (Applic	able to all LRF	ts, unless otherwise noted.)		Indicators for Problematic Hydric Soils ³ :
Histosol ((A1)		Sandy Gleyed Matrix ((S4)	1 cm Muck (A9) (LRR I, J)
	ipedon (A2)		Sandy Redox (S5)		Coast Prairie Redox (A16) (LRR F, G, H)
Black His			Stripped Matrix (S6)	~	Dark Surface (S7) (LRR G)
	n Sulfide (A4)		Loamy Mucky Mineral		High Plains Depressions (F16)
	Layers (A5) (LRR		Loamy Gleyed Matrix	(F2)	(LRR H outside of MLRA 72 & 73)
	ck (A9) (LRR F, G,		Depleted Matrix (F3)	-0)	Reduced Vertic (F18)
	Below Dark Surfac rk Surface (A12)	æ (A11)	Redox Dark Surface (Depleted Dark Surface)	23	Red Parent Material (TF2) Very Shallow Dark Surface (TF12)
	ucky Mineral (S1)		Redox Depressions (F		Other (Explain in Remarks)
	ucky Peat or Peat	(S2) (LRR G. H		Sector and the sector of the s	³ Indicators of hydrophytic vegetation and
	cky Peat or Peat (S		(MLRA 72 & 73 of	2 2	wetland hydrology must be present,
					unless disturbed or problematic.
Restrictive L	ayer (if present):				1 × 1
Type:					
Depth (inc	hes):				Hydric Soil Present? Yes No
Remarks:	the second s				
				1.1	
HYDROLOG	GY				-
Wetland Hyd	rology Indicators	2 D			12
Primary Indica	ators (minimum of	one required; ch	eck all that apply)		Secondary Indicators (minimum of two required)
	Vater (A1)		Salt Crust (B11)		Surface Soil Cracks (B6)
	er Table (A2)		Aquatic Invertebrates (B	13)	Sparsely Vegetated Concave Surface (B8)
Saturatio			Hydrogen Sulfide Odor (Drainage Patterns (B10)
Water Ma			Dry-Season Water Table	5	 Oxidized Rhizospheres on Living Roots (C3)
	t Deposits (B2)		Oxidized Rhizospheres of	. Sharaha ay marka ay marka	
	osits (B3)		(where not tilled)	in Living roots (c	Crayfish Burrows (C8)
	t or Crust (B4)		Presence of Reduced Iro	n (C4)	
Iron Depo	e 6		Thin Muck Surface (C7)		Geomorphic Position (D2)
	n Visible on Aerial	Imagery (B7)	Other (Explain in Remark	(8)	FAC-Neutral Test (D5)
	ained Leaves (B9)			,	Frost-Heave Hummocks (D7) (LRR F)
Field Observ					
			Depth (inches):		
Surface Wate			Deput (inches).		
Surface Wate	r Present?	Ves No	Depth (inches): 12+		
Water Table F	r Present?		Depth (inches): 12 +		
Water Table F Saturation Pre	r Present?) Present?) esent?)	17	Depth (inches): <u>2 +</u> Depth (inches): <u>2 +</u>	Wetla	nd Hydrology Present? Yes No
Water Table F Saturation Pre (includes capi	r Present?) Present?) esent?) illary fringe)	res No No	10 :		
Water Table F Saturation Pro (includes capi	r Present?) Present?) esent?) illary fringe)	res No No	Depth (inches): 12+		
Water Table F Saturation Pro (includes capi Describe Rec	r Present?) Present?) esent?) illary fringe)	res No No	Depth (inches): 12+		
Water Table F Saturation Pre (includes capi	r Present?) Present?) esent?) illary fringe)	res No No	Depth (inches): 12+		
Water Table F Saturation Pro (includes capi Describe Rec	r Present?) Present?) esent?) illary fringe)	res No No	Depth (inches): 12+		
Water Table F Saturation Pro (includes capi Describe Rec	r Present?) Present?) esent?) illary fringe)	res No No	Depth (inches): 12+		

Project/Site: Davis	_ City/County: <u>El Paso</u> Sampling Date: <u>5/23/23</u>
Applicant/Owner: Davis	State: CO _ Sampling Point: W9= UP
Investigator(s): Jan Davzvardis + Grant Gurnée	_ Section, Township, Range: <u>533;34, T125, R64W</u>
Landform (hillslope, terrace, etc.): <u>+evvace</u>	Local relief (concave, convex, none): Con Cave Slope (%): 1-9%
Subregion (LRR): Lat:	38,961083°N Long: -104, 543390°W Datum: WGS EU
Soil Map Unit Name: <u>B</u> Blakeland lamy sand	1 1-9% slopes NWI classification:/A
Are climatic / hydrologic conditions on the site typical for this time of	f year? Yes V No (If no, explain in Remarks.)
Are Vegetation, Soil, or Hydrology significan	ntly disturbed? Are "Normal Circumstances" present? Yes V No No
Are Vegetation, Soil, or Hydrology naturally	problematic? (If needed, explain any answers in Remarks.)
SUMMARY OF FINDINGS – Attach site map showing	ng sampling point locations, transects, important features, etc.
Hydrophytic Vegetation Present? Yes No	2

Hydrophytic Vegetation Present? Hydric Soil Present? Wetland Hydrology Present?	Yes Yes Yes	Is the Sampled Area within a Wetland?	Yes No	
Remarks:		e 17		
		<i>t</i>		
· ·				1

VEGETATION – Use scientific names of plants.

Tree Stratum (Plot size:) 1 2			Status	Dominance Test workshee Number of Dominant Specie That Are OBL, FACW, or FA (excluding FAC-):	es AC	(A)
2				Total Number of Dominant Species Across All Strata:	(B)
Sapling/Shrub Stratum (Plot size:) 1		= Total Cov	er _	Percent of Dominant Specie That Are OBL, FACW, or FA		A/B)
				Prevalence Index workshe	et:	
2				Total % Cover of.	Multiply by:	
3				OBL species	2011	
4				FACW species		
5				FAC species		
Herb Stratum (Plot size:)		= Total Cov	rer	FACU species		1
1. Boutploug gracilis	70	Y	UPL	UPL species		
2. Junaspatticus	15	Y	OBL	Column Totals:		(B)
3. Machaerantheva tanacetifolia		<u> </u>	VA	Prevalence Index [,] = B	/Å =	
4				Hydrophytic Vegetation In		
5				1 - Rapid Test for Hydro		- 1
6				2 - Dominance Test is >		
7				3 - Prevalence Index is		
8				4 - Morphological Adapt		orting
9			<u></u>	data in Remarks or c	on a separate sheet)	nung
10	-100-			Problematic Hydrophytic	c Vegetation ¹ (Explain)	
<u>Woody Vine Stratum</u> (Plot size:) 1		= Total Cov	ver	¹ Indicators of hydric soil and be present, unless disturbed		ust
2.				Hydrophytic	1	
% Bare Ground in Herb Stratum	100	= Total Cov	ver	Vegetation	No	
Remarks:						

US Army Corps of Engineers

C	0	1	
Э	U	ł	L

Sampling Point: M.9-UPL.

A CARLES AND A C		nfirm the absence of indicators.)
Depth <u>Matrix</u> (inches) Color (moist) %	Redox Features	2 Tostura Domodo
$\frac{11101000}{(2-10)}$	Color (moist) % Type ¹ Loc	1
<u>0-10 107R92</u>		Loam Dry
	· · · · · · · · · · · · · · · · · · ·	
¹ Type: C=Concentration, D=Depletion, RM	Reduced Matrix, CS=Covered or Coated San	d Grains. ² Location: PL=Pore Lining, M=Matrix.
Hydric Soil Indicators: (Applicable to a	I LRRs, unless otherwise noted.)	Indicators for Problematic Hydric Soils ³ :
Histosol (A1)	Sandy Gleyed Matrix (S4)	1 cm Muck (A9) (LRR I, J)
Histic Epipedon (A2)	Sandy Redox (S5)	Coast Prairie Redox (A16) (LRR F, G, H)
Black Histic (A3)	Stripped Matrix (S6)	Dark Surface (S7) (LRR G)
Hydrogen Sulfide (A4)	Loamy Mucky Mineral (F1)	High Plains Depressions (F16)
Stratified Layers (A5) (LRR F) 1 cm Muck (A9) (LRR F, G, H)	Loamy Gleyed Matrix (F2) Depleted Matrix (F3)	(LRR H outside of MLRA 72 & 73)
Depleted Below Dark Surface (A11)	Depleted Matrix (F3) Redox Dark Surface (F6)	Reduced Vertic (F18) Red Parent Material (TF2)
Thick Dark Surface (A12)	Depleted Dark Surface (F7)	Very Shallow Dark Surface (TF12)
Sandy Mucky Mineral (S1)	Redox Depressions (F8)	Other (Explain in Remarks)
2.5 cm Mucky Peat or Peat (S2) (LRR		³ Indicators of hydrophytic vegetation and
5 cm Mucky Peat or Peat (S3) (LRR F) (MLRA 72 & 73 of LRR H)	wetland hydrology must be present,
Dentil and the second		unless disturbed or problematic.
Restrictive Layer (if present):		1
Type:		
Depth (inches):		Hydric Soil Present? Yes No
Remarks:		
	14	
HYDROLOGY		-
Wetland Hydrology Indicators:		
Primary Indicators (minimum of one require	ed; check all that apply)	Secondary Indicators (minimum of two required)
Surface Water (A1)	Salt Crust (B11)	
LE-L More Table (10)		SUFFACE SOIL CLACKS (BD)
High Water Table (A2)	Aquatic Invertebrates (B13)	Surface Soil Cracks (B6) Sparsely Vegetated Concave Surface (B8)
High Water Table (A2) Saturation (A3)	Aquatic Invertebrates (B13)	Sparsely Vegetated Concave Surface (B8)
		Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10)
Saturation (A3)	Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1)	 Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3)
Saturation (A3) Water Marks (B1)	Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2)	 Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3)
Saturation (A3) Water Marks (B1) Sediment Deposits (B2)	 Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living Ro 	 Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled)
 Saturation (A3) Water Marks (B1) Sediment Deposits (B2) Drift Deposits (B3) Algal Mat or Crust (B4) Iron Deposits (B5) 	 Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living Ro (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7) 	 Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrøws (Cå)
 Saturation (A3) Water Marks (B1) Sediment Deposits (B2) Drift Deposits (B3) Algal Mat or Crust (B4) 	 Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living Ro (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7) 	 Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrews (C8) Saturation Visible on Aerial Imagery (C9)
 Saturation (A3) Water Marks (B1) Sediment Deposits (B2) Drift Deposits (B3) Algal Mat or Crust (B4) Iron Deposits (B5) 	 Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living Ro (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7) 	 Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrøws (C8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2)
 Saturation (A3) Water Marks (B1) Sediment Deposits (B2) Drift Deposits (B3) Algal Mat or Crust (B4) Iron Deposits (B5) Inundation Visible on Aerial Imagery (B 	 Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living Ro (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7) 	 Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrøws (C8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2) FAC-Neutral Test (D5)
 Saturation (A3) Water Marks (B1) Sediment Deposits (B2) Drift Deposits (B3) Algal Mat or Crust (B4) Iron Deposits (B5) Inundation Visible on Aerial Imagery (B) Water-Stained Leaves (B9) 	 Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living Ro (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7) Other (Explain in Remarks) 	 Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrøws (C8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2) FAC-Neutral Test (D5)
 Saturation (A3) Water Marks (B1) Sediment Deposits (B2) Drift Deposits (B3) Algal Mat or Crust (B4) Iron Deposits (B5) Inundation Visible on Aerial Imagery (B Water-Stained Leaves (B9) Field Observations: 	Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living Ro (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7) Thin Muck Surface (C7) Other (Explain in Remarks) No Depth (inches):	 Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrøws (C8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2) FAC-Neutral Test (D5)
Saturation (A3) Water Marks (B1) Sediment Deposits (B2) Drift Deposits (B3) Algal Mat or Crust (B4) Iron Deposits (B5) Iron Deposits (B5) Inundation Visible on Aerial Imagery (B Water-Stained Leaves (B9) Field Observations: Surface Water Present? Yes Water Table Present? Yes Saturation Present? Yes	Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living Ro (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7) Thin Muck Surface (C7) Other (Explain in Remarks) No Depth (inches): Depth (inches):	 Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrøws (C8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2) FAC-Neutral Test (D5)
Saturation (A3) Water Marks (B1) Sediment Deposits (B2) Drift Deposits (B3) Algal Mat or Crust (B4) Iron Deposits (B5) Iron Deposits (B5) I	Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living Ro (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7) Thin Muck Surface (C7) Other (Explain in Remarks) No Depth (inches): No Depth (inches): V	Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrøws (Č8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2) FAC-Neutral Test (D5) Frost-Heave Hummocks (D7) (LRR F) Vetland Hydrology Present? Yes No
Saturation (A3) Water Marks (B1) Sediment Deposits (B2) Drift Deposits (B3) Algal Mat or Crust (B4) Iron Deposits (B5) Iron Deposits (B5) I	Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living Ro (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7) Thin Muck Surface (C7) Other (Explain in Remarks) No Depth (inches): Depth (inches):	Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrøws (Č8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2) FAC-Neutral Test (D5) Frost-Heave Hummocks (D7) (LRR F) Vetland Hydrology Present? Yes No
Saturation (A3) Water Marks (B1) Sediment Deposits (B2) Drift Deposits (B3) Algal Mat or Crust (B4) Iron Deposits (B5) Iron Deposits (B5) Iron Deposits (B5) Inundation Visible on Aerial Imagery (B Water-Stained Leaves (B9) Field Observations: Surface Water Present? Yes Water Table Present? Yes Saturation Present? Yes (includes capillary fringe) Describe Recorded Data (stream gauge, m	Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living Ro (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7) Thin Muck Surface (C7) Other (Explain in Remarks) No Depth (inches): No Depth (inches): V	Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrøws (Č8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2) FAC-Neutral Test (D5) Frost-Heave Hummocks (D7) (LRR F) Vetland Hydrology Present? Yes No
Saturation (A3) Water Marks (B1) Sediment Deposits (B2) Drift Deposits (B3) Algal Mat or Crust (B4) Iron Deposits (B5) Iron Deposits (B5) I	Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living Ro (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7) Thin Muck Surface (C7) Other (Explain in Remarks) No Depth (inches): No Depth (inches): V	Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrøws (Č8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2) FAC-Neutral Test (D5) Frost-Heave Hummocks (D7) (LRR F) Vetland Hydrology Present? Yes No
Saturation (A3) Water Marks (B1) Sediment Deposits (B2) Drift Deposits (B3) Algal Mat or Crust (B4) Iron Deposits (B5) Iron Deposits (B5) Iron Deposits (B5) Inundation Visible on Aerial Imagery (B Water-Stained Leaves (B9) Field Observations: Surface Water Present? Yes Water Table Present? Yes Saturation Present? Yes (includes capillary fringe) Describe Recorded Data (stream gauge, m	Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living Ro (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7) Thin Muck Surface (C7) Other (Explain in Remarks) No Depth (inches): No Depth (inches): V	Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrøws (Č8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2) FAC-Neutral Test (D5) Frost-Heave Hummocks (D7) (LRR F) Vetland Hydrology Present? Yes No
Saturation (A3) Water Marks (B1) Sediment Deposits (B2) Drift Deposits (B3) Algal Mat or Crust (B4) Iron Deposits (B5) Inundation Visible on Aerial Imagery (B Water-Stained Leaves (B9) Field Observations: Surface Water Present? Yes Water Table Present? Yes Saturation Present? Yes (includes capillary fringe) Describe Recorded Data (stream gauge, m	Aquatic Invertebrates (B13) Hydrogen Sulfide Odor (C1) Dry-Season Water Table (C2) Oxidized Rhizospheres on Living Ro (where not tilled) Presence of Reduced Iron (C4) Thin Muck Surface (C7) Thin Muck Surface (C7) Other (Explain in Remarks) No Depth (inches): No Depth (inches): V	Sparsely Vegetated Concave Surface (B8) Drainage Patterns (B10) Oxidized Rhizospheres on Living Roots (C3) (where tilled) Crayfish Burrøws (Č8) Saturation Visible on Aerial Imagery (C9) Geomorphic Position (D2) FAC-Neutral Test (D5) Frost-Heave Hummocks (D7) (LRR F) Vetland Hydrology Present? Yes No