

Final

**Front Range Midway Solar Project
Wetlands, Waterbodies, and
Threatened, Endangered, and Species of Special Concern
Survey Report**

Prepared for:

Front Range Midway Solar, LLC

Prepared by:

Elizabeth Lack

Western EcoSystems Technology, Inc.
415 West 17th Street, Suite 200
Cheyenne, Wyoming 82001

August 7, 2015



TABLE OF CONTENTS

Introduction	1
Project Area Description	1
Methods	3
Wetlands and Waterbodies	3
Threatened, Endangered, and Species of Special Concern and Habitat	4
Results.....	5
Wetlands and Waterbodies	5
Threatened, Endangered, and Species of Special Concern and Habitat	7
Threatened and Endangered Species	7
State Species of Special Concern	9
Literature Cited	11

LIST OF TABLES

Table 1. Soils in the Project Area	3
Table 2. Federal and State Threatened, Endangered and Species of Special Concern	4

LIST OF FIGURES

Figure 1. Vicinity Map.....	2
Figure 2. Potential Wetland and Waterbody Features Investigated in the Field	6

LIST OF APPENDICES

APPENDIX A

Agency Letters

APPENDIX B

Photo Documentation of Potential Wetland and Waterbody Features Investigated at the Front Range Midway Solar Project Site

APPENDIX C

CDOW Recommended Survey Protocol and Actions to Protect Nesting Burrowing Owls

INTRODUCTION

Western EcoSystems Technology Inc. (WEST) conducted a survey at Front Range Midway Solar, LLC's¹ proposed project site to document any wetlands or other waterbodies that would be protected by the Clean Water Act and any potential occurrences or habitat for threatened or endangered species protected under the Endangered Species Act (ESA) or Colorado State law. The project site is located in El Paso County, Colorado, just west of Interstate 25 (I-25) and about 20 miles south of downtown Colorado Springs (Figure 1). The site will accommodate up to 100 megawatts (MW) of photovoltaic solar generating capacity and encompass approximately 1,085 acres of land. This survey was conducted to provide supporting information for compliance project environmental review, as well as compliance with the Clean Water Act and Endangered Species Act.

PROJECT AREA DESCRIPTION

The project area is located on the west side of I-25; a landfill is located to the south, a housing development consisting of 2.5-acre lots to the northwest, rangeland to the north, and a gravel pit adjacent to the site on the east. Other facilities nearby the project area include Pikes Peak International Raceway about 1.5 miles to the north and Fort Carson Military Reservation about one mile to the west. An electrical substation and the natural gas-fired Southwest Generation Power Plant are located within the project area (but are not included as part of the project area) at the west-central part of the site and several transmission lines connect to these facilities. Two fenced telecommunications compounds are also located within the project area.

The site is within Land Resource Region G, Western Great Plains (NRCS 2006). The project area is flat to gently rolling, at elevations ranging from approximately 5,360 to 5,520 feet. Surface runoff is generally to the east and flows to Fountain Creek, which flows to the south along the east side of I-25 to Pueblo where it joins the Arkansas River. The National Hydrology Dataset (NHD) portrays the surface water drainage network on maps; these are the blue lines seen on U.S. Geological Survey (USGS) topographic maps. No "blue lines" occur in the project area. The nearest named creek on a USGS topographic map is Sand Creek, over one-half mile south of the project area. The National Wetland Inventory (NWI) maps wetlands and deep water habitats of the U.S. According to the NWI, no wetlands occur at the project site.

Four soil map units are found in the project area; none are hydric soils. Table 1 summaries soils found in the project area.

¹ The project proponent, Front Range Midway Solar, LLC, is a wholly owned subsidiary of Tradewind Energy, LLC.




Figure 1. Vicinity Map

Table 1. Soils in the Project Area

Soil Map Unit	Soil Description
Kim loam, 1 to 8 percent slopes	Deep, well drained soils formed in calcareous loamy sediment on fans and uplands. Permeability is moderate.
Schamber-Razor complex, 8 to 50 percent slopes	Deep to moderately deep, well drained, gently rolling to steep soils on eroded breaks and remnants of granite outwash over shale. Permeability is slow to rapid.
Wilid silt loam, 0 to 3 percent slopes	Deep, well-drained soil formed in calcareous, silty eolian material. Permeability is moderate.
Fort loam, 1 to 5 percent slopes, cool	Deep, well drained soils formed from loamy eolian deposits on plains. Permeability is moderately high.

Source: Web Soil Survey, USDA Natural Resources Conservation Service

The natural vegetation of the project area is short-grass prairie. According to USGS National Land Cover Database, the primary cover type in the project area is grassland/herbaceous with a small area of scrub/shrub. The scrub/shrub classification includes areas dominated by shrubs less than five meters tall with a shrub canopy cover typically greater than 20 percent of total vegetation. This class includes true shrubs, young trees in an early successional stage, or trees stunted from environmental conditions. During a June 10, 2015 site visit, cane cholla (*Cylindropuntia imbricata*) was observed to be common throughout most of the grassland in the project area. Juniper (*Juniperus scopulorum*) trees were observed scattered in some of the drainage ways and at the northwest part of the project area.

METHODS

Wetlands and Waterbodies

Prior to conducting the field survey, a WEST biologist reviewed USGS topographic maps, Natural Resource Conservation Service (NRCS) web soil survey data, Google Earth aerial photography, and NWI data. Based on this review, all areas that could potentially be classified as a water of the U.S., including wetlands, were investigated in the field.

Two WEST biologists conducted the field survey on June 10, 2015. Wetland delineations followed the 1987 *Corps of Engineers Wetlands Delineation Manual* (Environmental Laboratory 1987) and the *Regional Supplement to the Corps of Engineers Wetland Delineation Manual: Great Plains Region (Version 2.0)* (USACE 2010). These manuals outline a three parameter approach for an area to be considered a wetland, in which all three parameters must be met. Hydrophytic vegetation must be the dominant vegetative cover, hydric soils must be present, and wetland hydrology must be present. The 2014 National Wetland Plant List (Lichvar et al. 2014) was used to determine the indicator status of plant species. Soil map units were determined from the NRCS Web Soil Survey website. The WEST biologists were to complete

wetland determination data forms for each sample point, and wetland boundaries were to be recorded on a Trimble 7X GPS unit with sub-foot accuracy.

Waterbodies were investigated in accordance with the Clean Water Rule. As such, the definition of “Waters of the United States” was taken from 40 CFR 230.3 (note: this is a prepublication version of the rule; the final rule was signed on 5/27/2015 and will become effective 60 days after publication in the Federal Register). Under this rule, tributaries must show physical features of flowing water (i.e., a bed, bank, and ordinary high water mark) to warrant protection under the Clean Water Act. All potential waterways were visited in the field to document the presence or absence of physical features of flowing water. The WEST biologists took photographs of to provide supporting documentation of the investigation.

Threatened, Endangered, and Species of Special Concern and Habitat

The project proponent previously completed an in-house Critical Issues Analysis, which included a list of federal and state threatened and endangered species in El Paso County (Table 2). The project proponent also sent letters to the U.S. Fish and Wildlife Service (Service) and Colorado Parks and Wildlife (CPW) requesting technical assistance review of the project. In their responses (Appendix A), the Service suggested an onsite habitat assessment for federally listed species and the CPW provided a list of state species of special concern in addition to threatened and endangered species. The species of special concern have been included in the evaluation (Table 2).

During the site visit on June 10, 2015, WEST biologists surveyed the project area to determine the habitat types present, and if any habitats might support listed threatened, endangered, and species of special concern. The survey was conducted by driving all roads in and around the project area and making observations. The substation properties in the middle of the project area were included in the visual evaluation. In addition, the Colorado Natural Heritage Program (CNHP) website was consulted to determine if any records of federal or state listed threatened or endangered species occur in the 7.5-minute quadrangle map (quad) in which the project is located (Buttes Quad).

Table 2. Federal and State Threatened, Endangered and Species of Special Concern – El Paso County, Colorado

Common Name	Scientific Name	State Status	Federal Status
Mexican Spotted Owl	<i>Strix occidentalis lucida</i>	Endangered	Threatened
Arkansas Darter	<i>Etheostoma cragini</i>	Threatened	Candidate Threatened
Greenback Cutthroat Trout	<i>Oncorhynchus clarki stomias</i>	Threatened	Threatened
Ute Ladies'-Tresses	<i>Spiranthes diluvialis</i>	None	Threatened
Pawnee Montane Skipper	<i>Hesperia leonardus montana</i>	None	Threatened
Black-Footed Ferret	<i>Mustela nigripes</i>	Endangered	EXP*

Common Name	Scientific Name	State Status	Federal Status
North American Wolverine	<i>Gulo gulo luscus</i>	Endangered	Proposed Threatened
Preble's Meadow Jumping Mouse	<i>Zapus hudsonius preblei</i>	Threatened	Threatened
Least Tern	<i>Sterna antillarum</i>	Endangered	Endangered
Piping Plover	<i>Charadrius melodus</i>	Threatened	Threatened
Whooping Crane	<i>Grus americana</i>	Endangered	Endangered
Pallid Sturgeon	<i>Scaphirhynchus albus</i>	None	Endangered
Plains Sharp-Tailed Grouse	<i>Tympanuchus phasianellus jamesii</i>	Endangered	None
Burrowing Owl	<i>Athene cunicularia</i>	Threatened	None
Lesser Prairie-Chicken	<i>Tympanuchus pallidicinctus</i>	Threatened	Proposed Threatened
River Otter	<i>Lontra canadensis</i>	Threatened	None
Prairie Dog	<i>Cynomys</i> spp.	Species of Special Concern	None (black-tailed prairie dog)
Swift Fox	<i>Vulpes velox</i>	Species of Special Concern	None
Mountain Plover	<i>Charadrius montanus</i>	Species of Special Concern	None
Townsend's Big-eared Bat	<i>Corynorhinus townsendii</i>	Species of Special Concern	None
Northern Leopard Frog	<i>Lithobates pipiens</i>	Species of Special Concern	None

Source: Critical Issues Analysis; Tradewind Energy, Inc., CPW letter to Tradewind Energy

RESULTS

Wetlands and Waterbodies

No wetlands occur in the project area. The WEST biologists investigated all areas that could potentially support wetlands and confirmed that no wetlands were found in the project area (Figure 2).




Figure 2. Potential Wetland and Waterbody Features Investigated in the Field

Other potential waterbodies that might be waters of the U.S. were also investigated in the field based on the pre-field data review. The investigation included areas that, topographically, could drain water (Figure 2). None of the drainage ways had physical features of flowing water, such as a bed, bank, or ordinary high water mark; therefore, they do not meet the definition of tributary and did not include characteristics of jurisdictional waters of the U.S. under the Clean Water Rule (see photographs, Appendix B). A dam occurs on one of the drainage ways, creating a stock pond (Photo 1, Appendix B). This stock pond had water at the time of the field investigation, probably due to timing of the survey in early June in a year with higher than average precipitation recorded for the month of May (NOAA 2015). The water appeared to be receding and likely dries up in late summer and in dry years in general. The banks were muddy and no hydrophytes were found along the bank, indicating water does not persist long enough or frequent enough to support hydrophytic vegetation. The stock pond did not include characteristics of a jurisdictional water of the U.S. (i.e., the Clean Water Rule specifically describes that artificial, constructed lakes and ponds constructed in dry land such as farm and stock watering ponds are not jurisdictional waters of the U.S.; Clean Water Rule Text § 230.3(s)(2)(iv)(B)).

Threatened, Endangered, and Species of Special Concern and Habitat

Threatened and Endangered Species

The list of federal and state threatened and endangered species in El Paso County prepared for a Critical Issues Analysis for the project included three fish (the state threatened Arkansas darter, the federal and state threatened greenback cutthroat trout, and the federal endangered pallid sturgeon). The field visit confirmed there are no waterbodies present at the project site that could support these fish species; therefore, these species could not occur there and the project would not affect these species. Similarly, no aquatic habitat is present at the project site for the state threatened river otter, so this species could not occur there and the project would not affect river otter.

Three other mammals were on the list of federal and state threatened and endangered species in El Paso County according to the Critical Issues Analysis prepared for the project: the federal and state endangered black-footed ferret, the state endangered North American wolverine, and the federal and state threatened Preble's meadow jumping mouse. The Service, in coordination with CPW (formerly the Colorado Division of Wildlife), has block-cleared all black-tailed prairie dog habitat in eastern Colorado, including El Paso County (USFWS 2009). This means the county has been determined to no longer contain any wild, free-ranging black-footed ferrets. Block clearance also means that the removal of black-tailed prairie dogs or their habitat (which provide habitat for black-footed ferrets) will no longer be required to meet the Service's survey guidelines for black-footed ferrets, or undergo consultation under Section 7 of the ESA (USFWS 2009). Based on the block clearance of El Paso County, the project would not affect the black-footed ferret. The North American wolverine occurs primarily in forested habitat and tundra. Because the project area does not contain habitat for this species, the project would not affect North American wolverine. The Preble's meadow jumping mouse inhabits well developed

riparian habitat with adjacent, relatively undisturbed grassland communities and a nearby water source. Well-developed riparian habitat includes a dense combination of grasses, forbs, and shrubs. No riparian habitat occurs in the project area and the only water source on the site is a seasonal stock pond with no shrubs in the riparian zone. Because habitat is not present at the project site for Preble's meadow jumping mouse, the project would not affect this species.

The list of federal and state threatened and endangered species in El Paso County included one butterfly, the federal threatened Pawnee montane skipper. This species has restricted range in portions of neighboring and nearby counties including Jefferson, Douglas, Teller and Park counties. This butterfly only occurs along the South Platte Canyon River drainage system in Ponderosa pine woodlands on moderately steep, granitic slopes. Because habitat does not occur for the Pawnee montane skipper in the project area and the project would not affect its habitat in nearby counties, the project would have no effect on this species.

Seven birds were on the list of federal and state threatened and endangered species in El Paso County: the state endangered and federal threatened Mexican spotted owl, the federal and state endangered least tern, the federal and state threatened piping plover, the federal and state endangered whooping crane, the state endangered plains sharp-tailed grouse, the state threatened burrowing owl, and the state and federal threatened lesser prairie chicken. Of these, the Mexican spotted owl would not be affected by the project because its habitat (forested mountains and canyons) is not present in the project area and this species would not occur there. Nesting habitat is not present in the project area for the least tern, piping plover, or whooping crane; however, because these species are migratory it is possible individuals could fly over the project area during migration. Even if this were to occur, the project is unlikely to affect these species because there is little to attract a migrating bird to the site (e.g., water), nor would solar panels (up to 10 feet in height) present a substantial collision hazard to migrating birds. Both the plains sharp-tailed grouse and the lesser prairie chicken are known from eastern Colorado; the plains sharp-tailed grouse to the northeast of the project area and the lesser prairie chicken to the southeast. While both are grassland species, both species' preferred habitat typically includes more shrubs than occur in the project area such as scrub oak and sand sage. The project is unlikely to affect these species because preferred shrub species are not present.

Burrowing owl nesting habitat consists of open areas with mammal burrows, such as the black-tailed prairie dog burrows that occur at the project site. Black-tailed prairie dogs were observed during the site visit and are active. Burrowing owls could occur and nest at the project site and be affected by the project. The CPW (formerly Colorado Division of Wildlife) has recommended survey protocols and actions to protect nesting burrowing owls (Appendix C). The protocol advises surveys for any activities occurring between March 15th and October 31 (burrowing owls are migratory and not expected to be present from November 1st to March 14th). Surveys are conducted in early mornings and evenings when the birds are most active and are conducted from a point with an unobstructed view of the prairie dog town. Multiple visits should be conducted to maximize the likelihood of detecting owls, if present. If owls are detected, CDOW recommends waiting to initiate activities until after November 1st or until it can be confirmed that

owls have left the prairie dog town, or carefully monitor the owls, noting and marking which burrows they are using. When all active burrows have been located and marked, activity can proceed in areas greater than 150 feet from the burrows with little danger to owls.

The list of federal and state threatened and endangered species in El Paso County included one plant species, the Ute ladies'-tresses orchid (*Spiranthes diluvialis*). This species occurs in moist meadows with perennial stream terraces, floodplains, oxbows, seasonally flooded river terraces, subirrigated or spring-fed abandoned stream channels and valleys, lakeshores, and human-modified wetlands. The on-site wetland and waterbody survey confirmed that habitat is not present in the project area for Ute ladies'-tresses orchid; therefore, the project would not affect this species.

State Species of Special Concern

The WEST biologists observed black-tailed prairie dogs in the north-central portion of the project area, east of the existing substation. Since prairie dogs are known to occur in the project area, the project will affect this species. The CPW recommends that prairie dogs be either moved alive to another location or humanely killed before any earth-moving occurs (Appendix A). CPW also recommends that since burrowing owls use prairie dog holes, the following should be observed:

- If construction is to occur between March 1 and October 31, the area should be surveyed for the presence of burrowing owls prior to any earth-moving taking place. The owls are susceptible to being buried and killed in their holes by construction activity. They are protected by law and killing one is illegal.
- If construction is to occur between November 1 and February 28, it is very unlikely that burrowing owls would be present since they migrate out of the state during winter (Appendix A).

Swift fox occurs on the shortgrass prairies of eastern Colorado and other central plains states (NatureServe 2015). Home range size ranges from a few hundred to a few thousand hectares (NatureServe 2015). They den in burrows, including prairie dog burrows. Most litters are born in March or early April and pups usually emerge by June 1 (NatureServe 2015). Because the project area includes habitat suitable for swift fox, including potential denning habitat, the project has potential to affect swift fox if they occur in the project area at the time of construction. If prairie dogs are removed prior to project-related earth-moving and outside of burrowing owl nesting season (March 1 through October 31), denning habitat for swift fox would also be eliminated outside of denning season when pups would be present, minimizing impacts to swift fox pups. The project would eliminate up to 1,085 acres of swift fox general habitat if the entire site is developed.

Mountain plover nest on high plains/shortgrass prairie habitat, including prairie dog towns in some areas (NatureServe 2015). In Colorado, nesting often occurs in shortgrass prairie with a history of heavy grazing or in low shrub semideserts. Nesting areas are characterized by very short vegetation, significant areas of bare ground (generally at least 30 percent bare ground),

and flat or gentle slopes (NatureServe 2015). Nesting begins in late April, incubation lasts for 29 days, and nestlings fledge in about 33 to 34 days. The project area includes some potential habitat suitable for mountain plover, including nesting habitat, particularly around the prairie dog burrows; however, vegetative cover appeared to be greater than 70 percent over most of the site based on observations during the site visit and vegetative appeared relatively tall for shortgrass prairie due to presence of cane cholla (up to several feet in height), which is common throughout the project site, as well as scattered trees in parts of the project area. The project would eliminate up to 1,085 acres of potential mountain plover habitat if the entire site is developed; however, most of the project area is not high quality habitat due to vegetative cover and structure.

Roosting habitat for Townsend's big-eared bat consists of spacious cavern-like structures such as caves and mines (Gruver and Keinath 2003). They forage along edge habitats (e.g., forested edges and intermittent streams), in forested habitat and along heavily vegetated stream corridors, and in open areas near wooded habitat though they appear to avoid open, grazed pasture land (Pierson et al. 1999). Water sources for drinking are open and accessible. Although roosting habitat is not present for Townsend's big-eared bat in the project area, the CPW report that a colony of Townsend's big-eared bat is located within a five-mile radius of the project area and bats might use the stock pond in the project area to drink and hunt insects (Appendix A). The stock pond would remain with development of the project and Townsend's big-eared bat could use it for foraging and water. The project would have little impact on the colony of Townsend's big-eared bat located within a five-mile radius because roosting habitat is not present in the project area and would not be affected, and the stock pond would continue to provide potential foraging opportunities and a water source for drinking.

Northern leopard frog live in the vicinity of springs, slow streams, marshes, bogs, ponds, canals, flood plains, reservoirs, and lakes (NatureServe 2015). They are usually in or near permanent water with rooted aquatic vegetation. In summer, they commonly inhabit wet meadows and fields, wintering sites are usually underwater (NatureServe 2015). Potential northern leopard frog habitat in the project area is limited to the stock pond. The WEST biologists observed water in the stock pond at the time of the field visit, probably due to timing in early June in a year with higher than average precipitation recorded for the month of May (NOAA 2015). The water appeared to be receding and likely dries up in late summer and in dry years in general. The banks were muddy and no hydrophytes were found along the bank, indicating water does not persist long enough or frequent enough to support hydrophytic or aquatic vegetation. No wet meadows or fields occur near the stock pond. The project area stock pond does not have preferred habitat features for the northern leopard frog, such as permanent water and rooted aquatic vegetation; the northern leopard frog is unlikely to occur there and project is unlikely to affect this species.

LITERATURE CITED

- Environmental Laboratory. 1987. *Corps of Engineers Wetlands Delineation Manual*. Technical Report Y-87-1. Department of the Army, Waterways Experiment Station, Corps of Engineers, Vicksburg, Mississippi.
- Gruver, J.C. and D.A. Keinath. Species Assessment for Townsend's Big-eared Bat (*Corynorhinus [=Plecotus] townsendii*) in Wyoming. Prepared for United States Department of the Interior Bureau of Land Management Wyoming State Office, Cheyenne, WY.
- Lichvar, R.W., M. Butterwick, N.C. Melvin, and W.N. Kirchner. 2014. *The National Wetland Plant List: 2014 Update of Wetland Ratings*. Phytoneuron 2014-41: 1-42.
- National Oceanic and Atmospheric Administration (NOAA). 2015. Climate Monitoring, National Overview – May 2015. Available at <https://www.ncdc.noaa.gov/sotc/national/201505>.
- Natural Resources Conservation Service (NRCS) Soil Survey Staff. 2006. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin. Agriculture Handbook 296 digital maps and attributes. Available online. Accessed July 2015.
- NatureServe. 2015. NatureServe Explorer: An Online Encyclopedia of Life [web application]. Version 7.1 NatureServe, Arlington, Virginia. Available <http://explorer.natureserve.org>. Accessed July 13, 2015.
- Pierson, E.D., M.C. Wackenhut, J.S. Altenbach, P. Bradley, P. Call, D. Genter, C.E. Harris, B.L. Keller, B. Lengus, L. Lewis, B. Luce, K.W. Navo, J.M. Perkins, S. Smith, and L. Welch. 1999. Species Conservation Assessment and Strategy for Townsend's Big-eared Bat (*Corynorhinus townsendii townsendii* and *Corynorhinus townsendii pallescens*). Idaho Department of Fish and Game, Boise, ID. In Gruver and Keinath 2003.
- U.S. Army Corps of Engineers. 2010. *Regional Supplement to the Corps of Engineers Wetland Delineation Manual: Great Plains Region* (Version 2.0), ed. J.S. Wakeley, R. W. Lichvar, and C.V. Noble. ERDC/EL TR-08-28. Vicksburg, MS: U.S. Army Engineer Research and Development Center.
- U.S. Fish and Wildlife Service, Mountain-Prairie Region. 2009. September 21, 2009 News Release: U.S. Fish and Wildlife Service and Colorado Department of Wildlife block clear eastern Colorado's black-tailed prairie dog habitat. Available online at: <http://www.fws.gov/mountain-prairie/pressrel/09-60.htm>

U.S. Fish and Wildlife Service. 2011. National Wetlands Inventory: Wetlands Online Mapper.
Available at: <http://www.fws.gov/wetlands/Data/GoogleEarth.html>.

APPENDIX A

Agency Letters

U.S. Fish and Wildlife Service

Colorado Parks and Wildlife



United States Department of the Interior



FISH AND WILDLIFE SERVICE
Ecological Services
Colorado Field Office
P.O. Box 25486, DFC (65412)
Denver, Colorado 80225-0486

IN REPLY REFER TO:

ES/CO: Solar Energy / El Paso County / Front Range-Midway Solar Project
TAILS: 06E24000-2014-TA-0805

JUL 29 2014

Ida Kitchen-Greenwell
Trade Wind Energy Inc.
16150 West 113th Street suite 105
Lenexa, KS 66219

Dear Ms. Kitchen-Greenwell:

Thank you for your email and letter to the U.S. Fish and Wildlife Service (Service) received July 16, 2014, regarding Trade Wind Energy's proposed Front Range-Midway solar photovoltaic (PV) project (project) located west of I-25 about 20 miles south of downtown Colorado Springs in El Paso County, Colorado.

Trade Wind Energy proposes to install and operate a solar array with capable of generating up to 100 MW of solar capacity on approximately 800 acres of vacant land, which is currently surrounded by infrastructure, including a regional landfill and a large electrical substation near the town of Fountain.

The PV panels will be affixed to a ground-mounted racking system supported by steel pylons driven into the ground. Light duty gravel service roads will be constructed within the solar array to provide access for ongoing maintenance. The solar array will be approximately 3 feet off ground surface and 1- feet in height, and will cover approximately 80% of the project area.

In preparation for a NEPA process and development of an Environmental Assessment you evaluated potential for threatened and endangered species to occur within the project area. Your report recommends that a habitat assessment be conducted to determine with greater certainty whether any T or E species habitats may be present in the area.

In response to your letter, we provide the following comments regarding:

1. Federally listed species;
2. Migratory birds;
3. Electrical transmission and distribution lines; and
4. State species of special concern, specifically the Gunnison's prairie dog.

The Service provides recommendations for threatened and endangered species under the authority of the Endangered Species Act of 1973 (ESA), as amended (16 U.S.C. 1531 *et seq.*).

Protective measures for migratory birds are provided under the authority of the Migratory Bird Treaty Act of 1918 (MBTA), as amended (16 U.S.C. 703 *et seq.*), and the Bald and Golden Eagle Protection Act of 1940 (BGEPA), as amended (16 U.S.C. 668 *et seq.*). We consider other fish and wildlife resources under the Fish and Wildlife Coordination Act (16 U.S.C. 661 *et seq.*) and the Fish and Wildlife Act (16 U.S.C. 742 *et seq.*).

1. Federally Listed Species

The proposed project is located within a developed area near the town of Fountain, Colorado. The primary vegetation type is grassland/herbaceous, which correlates to short-grass prairie with some scrub/shrub land cover. Given the land cover types, we would agree that an on-site habitat assessment should be conducted.

2. Migratory Birds and Bald and Golden Eagles

Activities associated with solar energy projects often include the removal of vegetation, underground burrows, or other structures used by migratory birds and eagles for nesting, roosting, perching, or foraging. During operation, solar energy facilities and their transmission lines may impact migratory birds by interrupting movements or by killing birds during collisions. Disturbed agricultural areas often provide foraging or ground nesting habitats for several migratory birds, such as the mountain plover (*Charadrius montanus*), and their conversion to solar farms may reduce or fragment available habitats. Therefore, we highlight the relevance of the MBTA and BGEPA to your project and provide recommendations intended to limit your project's impacts on migratory birds and eagles.

The Migratory Bird Treaty Act (MBTA):

The MBTA protects migratory birds, nests, and eggs from possession, sale, purchase, barter, transport, import, export, and take. Under the MBTA, it is unlawful unless permitted by regulations to pursue, hunt, take, capture, kill, or attempt to pursue, hunt, take, capture, or kill any migratory birds by any means or in any manner. The MBTA applies to 1,007 species of migratory birds identified in 50 CFR. § 10.13 and "take" is defined in 50 CFR § 10.12. The MBTA does not require intent to be proven, there is no incidental take statement, and the ESA does not absolve individuals or companies from liability under the MBTA. Unless permitted by the Service, the MBTA prohibits any intentional or unintentional activity that results in the take of migratory birds. Although the MBTA does not protect the habitats of migratory birds, activities that affect habitats and result in take of migratory birds do violate the MBTA.

The Bald and Golden Eagle Protection Act (BGEPA):

The BGEPA prohibits individuals and companies from knowingly, or with wanton disregard for the consequences of the Act, taking any bald or golden eagles or their body parts, nests, chicks, or eggs, which includes collection, molestation, disturbance, or killing. The BGEPA affords eagles additional protections beyond those provided by the MBTA by making it unlawful to "disturb" eagles. "Disturb" means to agitate or bother a bald or golden eagle to a degree that causes, or is likely to cause, injury to an eagle or decreases its productivity or results in nest abandonment due to interference with breeding, feeding, or sheltering behaviors. A permitting

process provides limited exceptions to the BGEPA's prohibitions and the Service has issued regulations concerning the permit procedures in 50 CFR Part 22.

Removing nests, destroying nests, or causing nest abandonment may constitute a violation of the MBTA and BGEPA. Removal of any active migratory bird nest or nest tree is prohibited. For golden eagles, permits for inactive nests are restricted to activities involving resource extraction for human health and safety. No permits will be issued for any active nest of any migratory bird species, unless removal of the active nest is necessary for reasons of human health and safety. Therefore, if nesting migratory birds are present within or near the project area, timing of activities is a significant consideration and should be addressed in the early phases of project planning. Nest manipulation is not allowed without a permit. If a permit cannot be issued, your project may need to be modified to ensure that take of any migratory bird, eagle, young, eggs, or nests will not occur.

Recommendations for migratory birds and eagles:

To minimize impacts to migratory birds, the Service recommends that construction occur outside the typical breeding season for migratory birds. Although the provisions of the MBTA apply year-round, most nesting activity occurs between April 1 and July 15. However, some migratory birds nest outside of this loosely defined period. If proposed activities must occur during the nesting season, or at any other time that may result in the take of migratory birds or eagles, the Service recommends that qualified biologists conduct pre-work field surveys of the affected habitats or structures, during the nesting season, to verify the presence or absence of migratory birds and eagles. Contact the Service's Colorado Field Office for guidance if surveys identify birds or nests that may be affected by project activities.

Enclosed, please find a copy of Colorado Parks and Wildlife's "Recommended Buffer Zones and Seasonal Restrictions for Colorado Raptors" (2008). We recommend reviewing these guidelines and incorporating the seasonal and buffer restrictions into your project design to avoid and minimize impacts to raptors and other migratory birds protected by the MBTA.

While adoption of these recommendations is voluntary, we remind Trade Wind Energy that the MBTA and BGEPA prohibit the take of migratory birds and eagles unless permitted by regulations. As mandated by our trust responsibilities, we immediately notify the Service's Office of Law Enforcement of any incidents of take at energy facilities.

It is not possible to absolve individuals, companies, or agencies from liability under the MBTA or BGEPA, even if they implement the guidelines or similar protective measures at their facilities. However, the Service's Office of Law Enforcement focuses its resources on investigating and prosecuting individuals and companies that take migratory birds without identifying and implementing all reasonable prudent and effective measures to avoid that take. It remains the applicant's responsibility to minimize the effects of their projects on migratory birds and other resources. For more information on MBTA and BGEPA regulations and their relevance to your project, please contact Craig Hansen of the Colorado Field Office at (303) 236-4749.

3. Electrical Transmission and Distribution Lines:

Solar energy facilities often require the development of new transmission and distribution lines. Overhead electrical lines concern the Service because published studies indicate that power lines can negatively affect wildlife. Collisions with power lines, power poles, and associated infrastructure often electrocute and kill birds, bats, and other wildlife. Projects may also permanently displace wildlife when activities alter or remove key components of important habitats. Early planning, coordination, and the strategic placement of power lines and associated facilities can avoid or reduce these impacts.

The Avian Power Line Interaction Committee (APLIC) developed guidelines and resources intended to address and mitigate electrocutions and collisions between wildlife and power lines. We recommend that you review and consider implementing these guidelines during the construction and operation of your electrical facilities. APLIC resources are available online at the following address:

<http://www.aplic.org/mission>

In Colorado, electrocutions at power lines are a serious threat to the ferruginous hawk (*Buteo regalis*), the golden eagle (*Aquila chrysaetos*), and other large raptors. In open prairies or agricultural fields, electrical poles often provide suitable perches or nest sites for birds of prey. As birds perch or build nests on power poles, their long wingspans easily touch electrical lines and complete circuits, effectively disrupting electrical service and often fatally electrocuting the bird. Undergrounding electrical lines eliminates the threat of electrocution and avian-caused power outages. Therefore, the Service recommends undergrounding electrical facilities whenever possible.

However, if undergrounding any overhead electrical line is not possible, we recommend that the proponents build overhead electrical lines with at least 10-foot cross arms on 3 phase lines, or at least 5 feet of spacing between electrical phases. Larger distances better accommodate long wingspans and may reduce electrocutions and power outages caused by birds at your power lines. APLIC provides additional recommendations to prevent electrocutions and power outages by discouraging perching and nesting.

4. State Species of Concern:

Our comments address federally listed species, federally designated critical habitats, and migratory birds. Please contact Colorado Parks and Wildlife (CPW) at (303) 297-1192 regarding any State species of special designation in Colorado that are not federally listed and that may occur within your project area. For example, the open areas within your project area may support colonies of the black-tailed prairie dog (*Cynomys ludovicianus*), a State species of special concern in Colorado.

The black-tailed prairie dog is a ground dwelling squirrel that lives in grasslands, including those in urban areas, disturbed right-of-ways, agricultural fields, and road or utility easements. Many grassland species, such as the burrowing owl (*Athene cunicularia*) depend on the underground burrows and colonies built by black-tailed prairie dogs. Due to their important value to the

prairie ecosystem and the many species that rely on them, we strongly encourage the conservation of prairie dogs.

To avoid and minimize impacts to prairie dogs or their dependent species, we recommend conducting preconstruction surveys for prairie dogs and their associated species. Design the project to avoid disturbing active colonies. If the project cannot avoid active colonies, relocate prairie dogs or consider donating them to a black-footed ferret or raptor recovery program. Contact CPW for more information on the regulations and guidelines that address the capture, transportation, and relocation of prairie dogs in Colorado.

The Service appreciates the opportunity to work with Trade Wind Energy on the proposed solar PV project. If we can be of any additional assistance, please contact the Colorado Field Office at 303-236-4773. Thank you for your concern endangered species and other natural resources.

Sincerely,



Susan C. Linner
Colorado Field Supervisor

Enclosure: CPW's recommended buffer zones and guidelines for raptors (2008)
Available online: <http://bit.ly/WXJYEh>



RECOMMENDED BUFFER ZONES AND SEASONAL RESTRICTIONS FOR COLORADO RAPTORS

Tolerance limits to disturbance vary among as well as within raptor species. As a general rule, Ferruginous Hawks and Golden Eagles respond to human activities at greater distances than do Ospreys and America Kestrels. Some individuals within a species also habituate and tolerate human activity at a proximity that would cause the majority of the group to abandon their nests. Other individuals become sensitized to repeated encroachment and react at greater distances. The tolerance of a particular pair may change when a mate is replaced with a less tolerant individual and this may cause the pair to react to activities that were previously ignored. Responses will also vary depending upon the reproductive stage. Although the level of stress is the same, the pair may be more secretive during egg laying and incubation and more demonstrative when the chicks hatch.

The term "disturbance" is ambiguous and experts disagree on what actually constitutes a disturbance. Reactions may be as subtle as elevated pulse rate or as obvious as vigorous defense or abandonment. Impacts of disturbance may not be immediately evident. A pair of raptors may respond to human intrusion by defending the nest, but well after the disturbance has passed, the male may remain in the vicinity for protection rather than forage to feed the nestlings. Golden eagles rarely defend their nests, but merely fly a half mile or more away and perch and watch. Chilling and over heating of eggs or chicks and starvation of nestlings can result from human activities that appeared not to have caused an immediate response.

A 'holistic' approach is recommended when protecting raptor habitats. While it is important for land managers to focus on protecting nest sites, equal attention should focus on defining important foraging areas that support the pair's nesting effort. Hunting habitats of many raptor species are extensive and may necessitate interagency cooperation to assure the continued nest occupancy. Unfortunately, basic knowledge of habitat use is lacking and may require documentation through telemetry investigations or intensive observation. Telemetry is expensive and may be disruptive so a more practical approach is to assume that current open space is important and should be protected.

Although there are exceptions, the buffer areas and seasonal restrictions suggested here reflect an informed opinion that if implemented, should assure that the majority of individuals within a species will continue to occupy the area. Additional factors, such as intervening terrain, vegetation screens, and the cumulative impacts of activities should be considered.

These guidelines were originally developed by CDOW raptor biologist Gerald R. Craig (retired) in December 2002. To provide additional clarity in guidance, incorporate new information, and update the conservation status of some species, the guidelines were revised in January 2008. Further revisions of this document may become necessary as additional information becomes available.

RECOMMENDED BUFFER ZONES AND SEASONAL RESTRICTIONS

BALD EAGLE

Nest Site:

No surface occupancy (beyond that which historically occurred in the area; see 'Definitions' below) within ¼ mile radius of active nests (see 'Definitions' below). Seasonal restriction to human encroachment (see 'Definitions' below) within ½ mile radius of active nests from October 15 through July 31. This closure is more extensive than the National Bald Eagle Management Guidelines (USFWS 2007) due to the generally open habitat used by Colorado's nesting bald eagles.

Winter Night Roost:

No human encroachment from November 15 through March 15 within ¼ mile radius of an active winter night roost (see 'Definitions' below) if there is no direct line of sight between the roost and the encroachment activities. No human encroachment from November 15 through March 15 within ½ mile radius of an active winter night roost if there is a direct line of sight between the roost and the encroachment activities. If periodic visits (such as oil well maintenance work) are required within the buffer zone after development, activity should be restricted to the period between 1000 and 1400 hours from November 15 to March 15.

Hunting Perch:

Diurnal hunting perches (see 'Definitions' below) associated with important foraging areas should also be protected from human encroachment. Preferred perches may be at varying distances from human encroachment and buffer areas will vary. Consult the Colorado Division of Wildlife for recommendations for specific hunting perches.

GOLDEN EAGLE

Nest Site:

No surface occupancy (beyond that which historically occurred in the area) within ¼ mile radius of active nests. Seasonal restriction to human encroachment within ½ mile radius of active nests from December 15 through July 15.

OSPREY

Nest Site:

No surface occupancy (beyond that which historically occurred in the area) within ¼ mile radius of active nests. Seasonal restriction to human encroachment within ¼ mile radius of active nests from April 1 through August 31. Some osprey populations have habituated and are tolerant to human activity in the immediate vicinity of their nests.

FERRUGINOUS HAWK

Nest Site:

No surface occupancy (beyond that which historically occurred in the area) within ½ mile radius of active nests. Seasonal restriction to human encroachment within ½ mile radius of active nests from February 1 through July 15. This species is especially prone to nest abandonment during incubation if disturbed.

RED-TAILED HAWK

Nest Site:

No surface occupancy (beyond that which historically occurred in the area) within 1/3 mile radius of active nests. Seasonal restriction to human encroachment within 1/3 mile radius of active nests from February 15 through July 15. Some members of this species have adapted to urbanization and may

tolerate human habitation to within 200 yards of their nest. Development that encroaches on rural sites is likely to cause abandonment.

SWAINSON'S HAWK

Nest Site:

No surface occupancy (beyond that which historically occurred in the area) within ¼ mile radius of active nests. Seasonal restriction to human encroachment within ¼ mile radius of active nests from April 1 through July 15. Some members of this species have adapted to urbanization and may tolerate human habitation to within 100 yards of their nest.

PEREGRINE FALCON

Nest Site:

No surface occupancy (beyond that which historically occurred in the area) within ½ mile radius of active nests. Seasonal restriction to human encroachment within ½ mile of the nest cliff(s) from March 15 to July 31. Due to propensity to relocate nest sites, sometimes up to ½ mile along cliff faces, it is more appropriate to designate 'Nesting Areas' that encompass the cliff system and a ½ mile buffer around the cliff complex.

PRAIRIE FALCON

Nest Site:

No surface occupancy (beyond that which historically occurred in the area) within ½ mile radius of active nests. Seasonal restriction to human encroachment within ½ mile radius of active nests from March 15 through July 15.

NORTHERN GOSHAWK

No surface occupancy (beyond that which historically occurred in the area) within ½ mile radius of active nests. Seasonal restriction to human encroachment within ½ mile radius of active nests from March 1 through September 15.

BURROWING OWL

Nest Site:

No human encroachment within 150 feet of the nest site from March 15 through October 31. Although Burrowing Owls may not be actively nesting during this entire period, they may be present at burrows up to a month before egg laying and several months after young have fledged. Therefore it is recommended that efforts to eradicate prairie dogs or destroy abandoned towns not occur between March 15 and October 31 when owls may be present. Because nesting Burrowing Owls may not be easily visible, it is recommended that targeted surveys be implemented to determine if burrows are occupied. More detailed recommendations are available in a document entitled "Recommended Survey Protocol and Actions to Protect Nesting Burrowing Owls" which is available from the Colorado Division of Wildlife

Recommended Buffer Zones and Seasonal Restrictions Around Raptor Use Sites

Species and Use	Buffer	Jan	Feb	Mar	Apr	May	June	July	Aug	Sept	Oct	Nov	Dec
Bald Eagle													
ACTIVE NEST - No Surface Occupancy	1/4 Mile												
ACTIVE NEST - No Human Encroachment	1/2 Mile												
ACTIVE WINTER NIGHT ROOST without a direct line of sight- No Human Encroachment	1/4 Mile												
ACTIVE WINTER NIGHT ROOST with a direct line of sight - No Human Encroachment	1/2 Mile												
HUNTING PERCH - No Human Encroachment	Contact CDOW												
Golden Eagle													
ACTIVE NEST - No Surface Occupancy	1/4 Mile												
ACTIVE NEST - No Human Encroachment	1/2 Mile												
Osprey													
ACTIVE NEST - No Surface Occupancy	1/4 Mile												
ACTIVE NEST - No Human Encroachment	1/2 Mile												
Ferruginous Hawk													
ACTIVE NEST - No Surface Occupancy	1/2 Mile												
ACTIVE NEST - No Human Encroachment	1/2 Mile												
Red-tailed Hawk													
ACTIVE NEST - No Surface Occupancy	1/3 Mile												
ACTIVE NEST - No Human Encroachment	1/3 Mile												
Swainson's Hawk													
ACTIVE NEST - No Surface Occupancy	1/4 Mile												
ACTIVE NEST - No Human Encroachment	1/4 Mile												
Peregrine Falcon													
ACTIVE NEST - No Surface Occupancy	1/2 Mile												
ACTIVE NEST - No Human Encroachment	1/2 Mile												
Prairie Falcon													
ACTIVE NEST - No Surface Occupancy	1/2 Mile												
ACTIVE NEST - No Human Encroachment	1/2 Mile												
Northern Goshawk													
ACTIVE NEST - No Surface Occupancy	1/2 Mile												
ACTIVE NEST - No Human Encroachment	1/2 Mile												
Burrowing Owl													
ACTIVE NEST - No Human Encroachment	150 feet												

= time period for which seasonal restrictions are in place.

DEFINITIONS

Active nest – Any nest that is frequented or occupied by a raptor during the breeding season, or which has been active in any of the five previous breeding seasons. Many raptors use alternate nests in various years. Thus, a nest may be active even if it is not occupied in a given year.

Active winter night roost – Areas where Bald Eagles gather and perch overnight, and sometimes during the day in the event of inclement weather. Communal roost sites are usually in large trees (live or dead) that are relatively sheltered from wind and are generally in close proximity to foraging areas. These roosts may also serve a social purpose for pair bond formation and communication among eagles. Many roost sites are used year after year.

Human encroachment – Any activity that brings humans in the area. Examples include driving, facilities maintenance, boating, trail access (e.g., hiking, biking), etc.

Hunting perch – Any structure on which a raptor perches for the purpose of hunting for prey. Hunting perches provide a view of suitable foraging habitat. Trees are often used as hunting perches, but other structures may also be used (utility poles, buildings, etc.).

Surface occupancy – Any physical object that is intended to remain on the landscape permanently or for a significant amount of time. Examples include houses, oil and gas wells, tanks, wind turbines, roads, tracks, etc.

CONTACT

For further information contact:

David Klute
Bird Conservation Coordinator
Colorado Division of Wildlife
6060 Broadway
Denver, CO 80216
Phone: 303-291-7320
Email: david.klute@state.co.us

REFERENCES

- Bechard, M.J., and J.K. Schmutz. 1995. Ferruginous Hawk (*Buteo regalis*). The Birds of North America Online (A. Poole, Ed.). Ithaca: Cornell Lab of Ornithology; Retrieved from the Birds of North America Online: <http://bna.birds.cornell.edu/bna/species/172>
- Buehler, D.A. 2000. Bald Eagle (*Haliaeetus leucocephalus*). The Birds of North America Online (A. Poole, Ed.). Ithaca: Cornell Lab of Ornithology; Retrieved from the Birds of North America Online: <http://bna.birds.cornell.edu/bna/species/506>
- Call, M. 1979. Habitat management guides for birds of prey. Technical Note No.338, U.S. Bureau of Land Management, Denver Service Center, Denver, CO. 69pp

- Energy Research and Development Administration (ERDA). 1977. EIA for CUI Venture application for geothermal loan guarantee (Beryl and Lund, Utah). EIA/GE/77-8. Washington, D.C. 109pp.
- England, A.S., M.J. Bechard, and C.S. Houston. 1997. Swainson's Hawk (*Buteo swainsoni*), The Birds of North America Online (A. Poole, Ed.). Ithaca: Cornell Lab of Ornithology; Retrieved from the Birds of North America Online: <http://bna.birds.cornell.edu/bna/species/265>
- Greater Yellowstone Bald Eagle Working Group. 1996. Greater Yellowstone bald eagle management plan: 1995 update. Greater Yellowstone Bald Eagle Working Group, Wyoming Game & Fish Dept., Lander WY 82520. 47p
- Grier, J.W., F.J. Gramlich, J. Mattisson, J.E. Mathisen, J.V. Kussman, J.B. Elder, and N.F. Green. 1983. The bald eagle in the northern United States. *Bird Cons.* 144-66.
- Haug, E.A., B.A. Millsap, and M.S. Martell. 1993. Burrowing Owl (*Athene cunicularia*), The Birds of North America Online (A. Poole, Ed.). Ithaca: Cornell Lab of Ornithology; Retrieved from the Birds of North America Online: <http://bna.birds.cornell.edu/bna/species/061>
- Holmes, Tamara L. 1993. Behavioral responses of grassland raptors to human disturbance. MS Thesis. Colo. State Univ., Fort Collins. 62pp.
- Holthuijzen, A.M.A., W.G. Eastland, A.R. Ansell, M.N. Kochert, R.D. Williams, and L.S. Young. 1990. Effects of blasting on behavior and productivity of nesting prairie falcons. *Wildl. Soc. Bull.* 18:270-281.
- Kochert, M. N., K. Stecnhof, C. L. McIntyre, and E. H. Craig. 2002. Golden Eagle (*Aquila chrysaetos*), The Birds of North America Online (A. Poole, Ed.). Ithaca: Cornell Lab of Ornithology; Retrieved from the Birds of North America Online: <http://bna.birds.cornell.edu/bna/species/684>
- Martin, D.J. 1973. Selected aspects of burrowing owl ecology and behavior. *Condor* 75:446-456.
- Northern States Bald Eagle Recovery Team. 1983. Northern States Bald Eagle Recovery Plan. U.S. Fish and Wildlife Service. 75pp.
- Olanderff, R. R., and W.D. Zeedyk. 1978. Land management for the conservation of endangered birds. Pages 419-428 in S.A. Temple, ed. *Endangered birds*. University of Wisconsin Press, Madison, Wisconsin.
- Poole, A.F., R.O. Bierregaard, and M.S. Martell. 2002. Osprey (*Pandion haliaetus*), The Birds of North America Online (A. Poole, Ed.). Ithaca: Cornell Lab of Ornithology; Retrieved from the Birds of North America Online: <http://bna.birds.cornell.edu/bna/species/683>
- Preston, C.R., and R.D. Beane. 1993. Red-tailed Hawk (*Buteo jamaicensis*), The Birds of North America Online (A. Poole, Ed.). Ithaca: Cornell Lab of Ornithology; Retrieved from the Birds of North America Online: <http://bna.birds.cornell.edu/bna/species/052>

- Reynolds, R., R.T. Graham, H.M. Reiser. 1992. Management recommendations for the northern goshawk in the southwestern United States. Gen. Tech. Rep. RM-217. Fort Collins, CO. U.S. Dept of Agri., Forest Service, Rocky Mountain Forest and Range Experiment Station. 90pp.
- Richardson, C.T. and C.K. Miller. 1997. Recommendations for protecting raptors from human disturbance: a review. Wildl. Soc. Bull. 25(3):634-638.
- Rocky Mountain/Southwest Peregrine Falcon Recovery Team. 1984. American peregrine falcon Rocky Mountain/Southwest population recovery plan. U.S. Fish and Wildlife Serv. 105pp.
- Squires, J.R., S.H. Anderson, and R. Oakleaf. 1993. Home range size and habitat-use patterns of nesting prairie falcons near oil developments in northeastern Wyoming. J. Field Ornithol. 64:1-10.
- Steenhof, Karen. 1998. Prairie Falcon (*Falco mexicanus*), The Birds of North America Online (Poole, Ed.). Ithaca: Cornell Lab of Ornithology; Retrieved from the Birds of North America Online: <http://bna.birds.cornell.edu/bna/species/346>
- Squires, J.R., and R.T. Reynolds. 1997. Northern Goshawk (*Accipiter gentilis*), The Birds of North America Online (A. Poole, Ed.). Ithaca: Cornell Lab of Ornithology; Retrieved from the Birds of North America Online: <http://bna.birds.cornell.edu/bna/species/298>
- Suter, G.W. and J.L. Joness. 1981. Criteria for Golden Eagle, Ferruginous Hawk, and Prairie Falcon nest site protection. J. Raptor Res. 15(1):12-18.
- Swenson, J.E. 1979. Factors affecting status and reproduction of ospreys in Yellowstone National Park. J. Wildl. Manage. 43:595-601.
- Thomsen, L. 1971. Behavior and ecology of burrowing owls on the Oakland Municipal Airport. Condor 73:177-192.
- U.S. Fish and Wildlife Service. 2007. National Bald Eagle Management Guidelines. <http://www.fws.gov/migratorybirds/issues/BaldEagle/NationalBaldEagleManagementGuidelines.pdf>
- White, C.M., N.J. Clum, T.J. Cade, and W.G. Hunt. 2002. Peregrine Falcon (*Falco peregrinus*), The Birds of North America Online (A. Poole, Ed.). Ithaca: Cornell Lab of Ornithology; Retrieved from the Birds of North America Online: <http://bna.birds.cornell.edu/bna/species/660>

Revised 02/2008



COLORADO
Parks and Wildlife

Department of Natural Resources

Southeast Region
4255 Sinton Road
Colorado Springs, CO 80907
P 719.227.5200 | F 719.227.5223

August 25, 2014

TradeWind Energy, Inc.
Jennifer A Dean
16150 West 113th Street Suite 105
Lenexa, KS 66219
jdean@tradewindenergy.com

Re: The Front Range-Midway Solar Project

Dear Ms. Dean,

Thank you for the opportunity to assist in the review of the desktop analysis which was done for The Front Range-Midway Solar Project (Project). The Project is expected to encompass approximately 800 acres in El Paso County, which will accommodate up to 100 MW of solar capacity.

The mission of Colorado Parks and Wildlife (CPW) is to perpetuate the wildlife resources of the state, to provide a quality state parks system, and to provide enjoyable and sustainable outdoor recreation opportunities that educate and inspire current and future generations to serve as active stewards of Colorado's natural resources. One of the ways we achieve our mission is to comment on land use proposals such as the request we received from you. Our goal is to provide complete, consistent and timely information to all entities who request comment on matters within our statutory authority and our mission.

CPW is generally very supportive of renewable energy products, provided that impacts to wildlife resources are considered and mitigated to the extent possible during the design, construction, and operation of the renewable energy facilities. District Wildlife Manager Cody Wigner has recently reviewed the documents associated with the Project and has visited the Project site. CPW believes the following species could be present on the Project site and has the following comments:

Threatened and Endangered Species:

Burrowing Owl - (State Threatened) Burrowing Owls may be present on the Project Site. Burrowing owls live and nest in prairie dog holes. Since prairie dogs were seen on the property, burrowing owls might use it during the summer.



Prairie Dogs - (Species of special concern) Are a high interest species, serve important ecological functions, and have been drastically reduced in numbers.

- CPW recommends that prairie dogs be either moved alive to another location or humanely killed before any earth-moving occurs; If killed, a properly licensed commercial applicator must be used and if relocated, a permit through the Division of Wildlife must be obtained; and

- Since burrowing owls use prairie dogs holes, the following should be observed:

- If construction is to occur between March 1 and October 31, the area should be surveyed for the presence of burrowing owls prior to any earth-moving taking place. The owls are susceptible to being buried and killed in their holes by construction activity. They are protected by law and killing one is illegal.

- If construction is to occur between November 1 and February 28, it is very unlikely that burrowing owls would be present since they migrate out of the state during the winter.

Swift Fox - (Species of special concern) The proposed development site is in the range of the Swift fox. It is a special concern species, because it was warranted as a federally threatened species, but precluded by other higher priority species, thus placing the species on the candidate list.

Mountain Plover - (Species of special concern) A grassland bird, are likely to be nesting in the Project area.

Townsend's Big Eared Bat - (Species of special concern) A colony of is located within a 5 mile radius of the Project area. The bats may frequent the small water hole in the Project area to drink and hunt insects.

Northern Leopard Frog - (Species of special concern) Possible habitat is located in the small wetland in the Project area.

For information on surveying for burrowing owls, please contact District Wildlife Manager Cody Wigner.

Habitat Loss:

CPW would also like TradeWind Energy to be conscious that long term habitat loss on the Project area will be a main impact on wildlife. Other wildlife species that can potentially be found on the Project site are: black tailed prairie dog, bobcat, cottontail rabbit, coyote, mule deer, white-tailed deer, elk, pronghorn, red fox, jack rabbit, mountain lion, skunks, variety of small burrowing rodents, a variety of reptiles which include snakes and lizards, and a variety of grassland birds, Golden eagle, Ferruginous hawk, Red-tailed hawk, Prairie falcon, and Swainson's hawk likely hunt nearby and within the prairie dog colony. These raptors may nest in the area. An annually active known Golden eagle nest is located within a 5-mile

radius of the project. CPW recommends that the wetland in the Project area remains undisturbed and contiguous with undeveloped land around it. The wetland provides possible habitat for the northern leopard frog, as well as a water source for all wildlife and possibly used by Townsend's big eared bats as hunting grounds.

For protected migratory bird species, CPW recommends maintaining buffer zones and seasonal restrictions (see attachment: Raptor Buffer Guidelines 2008).

CPW would also like to make sure TradeWind Energy is aware of the Colorado PUC Environmental Renewable Energy Standards for Electric Utilities. For a copy of these regulations, please contact District Wildlife Manager Cody Wigner.

Thank you again for the opportunity to assist in the review of the desktop analysis which was done for The Front Range-Midway Solar Project. Please do not hesitate to contact CPW about ways to continue to maximize wildlife value while minimizing potential conflicts on the Project. CPW appreciates having the opportunity to get involved early in the process of the Project. If you have further questions please contact District Wildlife Manager Cody Wigner at (719) 227-5287 or via email at cody.wigner@state.co.us.


Sincerely,



Frank McGee
Area Wildlife Manager

Cc: SE Region Files
Area 14 Files
C. Wigner, DWM

APPENDIX B
Photo Documentation of Potential Wetland and Waterbody Features Investigated at the
Front Range Midway Solar Project Site



Location and Direction of Photographs taken in Project Area



Photo 1. Stock pond (view from dam looking southwest)



Photo 2.

Drainage way at northcentral part of the project area, looking northeast from dam



Photo 3.
Drainage way at northcentral part of the project area, looking southwest (view from upper end of stock pond)



Photo 4.
Drainage way at southeast part of the project area, looking northwest



Photo 5.
Drainage way at southeast part of the project area, looking south



Photo 6.
Drainage way at southeast part of the project area, looking north from road along southern border of project area



Photo 7.

Drainage way at southwest part of the project area, looking northwest from road along southern border of project area



Photo 8.

Drainage way at southwest part of the project area, looking northwest (upstream from Photo 7 location)



Photo 9.

Drainage way at southwest part of the project area, looking south from road through center of project area



Photo 10.

Drainage way at southwest part of the project area, looking north from road through center of project area (no discernable drainage pattern on landscape)

APPENDIX C

CDOW Recommended Survey Protocol and Actions to Protect Nesting Burrowing Owls



RECOMMENDED SURVEY PROTOCOL AND ACTIONS TO PROTECT NESTING BURROWING OWLS

Western Burrowing Owls (*Athene cunicularia hypugaea*) are commonly found in prairie dog towns throughout Colorado. Burrowing owls require prairie dog or other suitable burrows (e.g. badger) for nesting and roosting. Burrowing owls are migratory, breeding throughout the western United States, southern Canada, and northern Mexico and wintering in the southern United States and throughout Mexico.

Federal and state laws prohibit the harming or killing of burrowing owls and the destruction of active nests. It is quite possible to inadvertently kill burrowing owls during prairie dog poisoning projects, removal of prairie dogs, destruction of burrows and prairie dogs using a concussive device, or during earth moving for construction. Because burrowing owls often hide in burrows when alarmed, it is not practical to haze the birds away from prairie dog towns prior to prairie dog poisoning/removal, burrow destruction, or construction activity. Because of this, the Colorado Division of Wildlife recommends surveying prairie dog towns for burrowing owl presence before potentially harmful activities are initiated.

The following guidelines are intended as advice on how to determine if burrowing owls are present in a prairie dog town, and what to do if burrowing owls are detected. These guidelines do not guarantee that burrowing owls will be detected if they are present. However, adherence to these guidelines will greatly increase the likelihood of detection.

Seasonal Timing

Burrowing owls typically arrive on breeding grounds in Colorado in late March or early April, with nesting beginning a few weeks later. Active nesting and fledging has been recorded and may be expected from late March through early August. Adults and young may remain at prairie dog towns until migrating to wintering grounds in late summer or early autumn.

Surveys should be conducted during times when burrowing owls may be present on prairie dog towns. Surveys should be conducted for any activities occurring between March 15th and October 31st. No burrowing owls are expected to be present between November 1st and March 14th.

Daily Timing

Burrowing owls are active throughout the day; however, peaks in activity in the morning and evening make these the best times for conducting surveys (Conway and Simon 2003). Surveys should be conducted in the early morning (1/2 hour before sunrise until 2 hours after sunrise) and early evening (2 hours before sunset until 1/2 hour after sunset).

Number and locations of survey points

Burrowing owls are most frequently located visually, thus, obtaining a clear view of the entire prairie dog town is necessary. For small prairie dog towns that can be adequately viewed in their entirety from a single location, only one survey point is necessary. The survey point should be selected to provide unobstructed views (with binoculars if necessary) of the entire prairie dog town

(burrow mounds and open areas between) and all nearby structures that may provide perches (e.g., fences, utility poles, etc.)

For prairie dog towns that can not be entirely viewed from a single location because of terrain or size, enough survey points should be established to provide unobstructed views of the entire prairie dog town and nearby structures that may provide perches. Survey locations should be separated by approximately 800 meters (1/2 mile), or as necessary to provide adequate visual coverage of the entire prairie dog town.

Number of surveys to conduct

Detection of burrowing owls can be highly variable and multiple visits to each site should be conducted to maximize the likelihood of detecting owls if they are present. At least three surveys should be conducted at each survey point. Surveys should be separated by approximately one week.

Conducting the survey

- **Weather Considerations** Because poor weather conditions may impact the ability to detect burrowing owls, surveys should only be conducted on days with little or no wind and no precipitation.
- **Passive surveys** Most burrowing owls are detected visually. At each survey location, the observer should *visually* scan the area to detect any owls that are present. Some burrowing owls may be detected by their call, so observers should also *listen* for burrowing owls while conducting the survey.

Burrowing owls are frequently detected soon after initiating a survey (Conway and Simon 2003). However, some burrowing owls may not be detected immediately because they are inconspicuous, are inside of burrows, or are not present on the site when the survey is initiated. We recommend that surveys be conducted for 10 minutes at each survey location.

- **Call-broadcast surveys** To increase the likelihood of detecting burrowing owls, if present, we recommend incorporating call-broadcast methods into burrowing owl surveys. Conway and Simon (2003) detected 22% more burrowing owls at point-count locations by broadcasting the primary male (*coo-coo*) and alarm (*quick-quick-quick*) calls during surveys. Although call-broadcast may increase the probability of detecting burrowing owls, most owls will still be detected visually.
- We recommend the following 10-minute timeline for incorporating call-broadcast methods (Conway and Simon 2003, C. Conway pers. commun.). The observer should scan the area for burrowing owls during the entire survey period.
 - 3 minutes of silence
 - 30 seconds call-broadcast of primary call (*coo-coo*)
 - 30 seconds silence
 - 30 seconds call-broadcast of primary call (*coo-coo*)
 - 30 seconds silence
 - 30 seconds call-broadcast of alarm call (*quick-quick-quick*)
 - 30 seconds silence
 - 4 minutes of silence

Calls can be broadcast from a "boom box", a portable CD or cassette player, or an mp3 player attached to amplified speakers. Calls should be broadcast loudly but without distortion.

Recordings of this survey sequence (compact disc or mp3 sent via email) are available free of charge by contacting:

David Klute
Bird Conservation Coordinator
Colorado Division of Wildlife
6060 Broadway
Denver, CO 80216
Phone: 303-291-7320
Email: David.Klute@state.co.us

Identification

Adult burrowing owls are small, approximately 9-11 inches. They are brown with white spotting and white barring on the chest. They have long legs in comparison to other owls and are frequently seen perching on prairie dog mounds or other suitable perches (e.g., fence posts, utility poles) near prairie dog towns. Juvenile burrowing owls are similar to adults but smaller, with a white/buff colored chest that lacks barring.

General information about burrowing owls is available from the Colorado Division of Wildlife website:

<http://wildlife.state.co.us/WildlifeSpecies/Profiles/Birds/BurrowingOwl.htm>

Additional identification tips and information are available from the U.S. Geological Survey Patuxent Wildlife Research Center website:

<http://www.mbr-pwrc.usgs.gov/id/framlst/i3780id.html>

What To Do If Burrowing Owls Are Present

If burrowing owls are confirmed to be present in a prairie dog town, there are two options before proceeding with planned activities:

1. Wait to initiate activities until after November 1st or until it can be confirmed that the owls have left the prairie dog town.
2. Carefully monitor the activities of the owls, noting and marking which burrows they are using. This is not easy to accomplish and will require considerable time, as the owls may use several burrows in a prairie dog town. When all active burrowing owl burrows have been located and marked, activity can proceed in areas greater than 150 feet from the burrows with little danger to the owls. Activity closer than 150 feet may endanger the owls.

Reference

Conway, C. J. and J. C. Simon. 2003. Comparison of detection probability associated with Burrowing Owl survey methods. *Journal of Wildlife Management* 67:501-511.

revised 02/2008

See also: "Controlling Prairie Dogs: Suggestions For Minimizing Risk To Non-Target Wildlife Species" Colorado Division of Wildlife 03/2007